

HEALTH INFORMATION designs

West Virginia Department of Health and Human Resources Bureau for Medical Services Drug Utilization Review Board September 20, 2017

OProfiles Reviewed	1,206)
Cases Identified	900	
OLetters Mailed		
O Prescribers	1,105	
O Pharmacies	971	
○ Responses		
O Prescribers	179	16.2%
 Pharmacies 	174	17.9%



April RDUR Criteria:

• Controlled substances

Orug-Drug Conflicts:

○ SSRI's & TCAs

Anticoagulants

O Benzodiazepines & buspirone

○ ACEI's & NSAIDs

ONSAIDs & Beta Blockers

Orug-Disease Interactions

 $\odot\,\text{NSAID}$ use in patients with diabetes



April Educational Intervention:

• Triple Drug Combination

- Patients on an opioid + skeletal muscle relaxant + benzodiazepine
- Education on risk of CNS depression, abuse/dependence, and diversion
- \odot Sent to all prescribers of at least one of the above for patients on the combination
 - o Cases Reviewed: 548
 - Letters Sent: 652



May RDUR Criteria:

Controlled substances

O Drug-Disease interactions

• Aripiprazole

- o Patients with seizure disorders
- o Patients with cardiovascular or cerebrovascular disease

O NSAIDS

- Patients with cardiovascular disease
- $\circ~$ Patients with renal impairment

O Drug-Drug Interactions:

- o NSAIDS /Warfarin
- $\,\circ\,$ Duplication of NSAIDS
- Cyclobenzaprine and TCAs
- $\,\circ\,$ SSRI's and other serotonergic agents
- Stimulants and other serotonergic agents



June RDUR Criteria:

Controlled substances

OUnderutilization

\odot Long-term as thma controllers

OTherapeutic duplication

\circ Antiulcer agents

- $\circ \text{Anxiolytics}$
- \odot Loop diuretics
- **O Skeletal Muscle Relaxants**

Clinical Appropriateness Polypsychotherapy



June Educational Intervention:

Identify new members on non-formulary test strips

Sent to prescriber's with patients on True Metrix
 Diabetic Products to inform affected patients

 $\,\circ\,$ Single letter to prescriber with all patients affected.

- \odot Letters sent included all potentially affected patients.
 - Cases Reviewed: 2,798
 - Letters Sent: 1,029



Proposed Educational Interventions

Opioid MME restriction for new members

 December mailed 1,297 MD letters for 4,735 affected members in current population

 Therapy with both a long-acting benzodiazepine and sedative/hypnotic

- MD letters to patients with concurrent or alternating claims
- **FDA's Drug Safety Communications**

 \odot Codeine and tramadol use in children



Distribution of Cases:

- Drug-Disease Interactions: 12%
 - Patients receiving a drug that may worsen or precipitate a medical condition.
- Drug-Drug Conflict: 61%
 - Patients receiving two or more drugs that may interact and produce unpredictable and undesirable effects.
- Over-utilization: 8%
 - Patients taking medications in apparently excessive doses or for excessive lengths of time.
- Non-compliance: 6%
 - Patients not taking medication according to directions, resulting in possible subtherapeutic response.
- Clinical Appropriateness: 12%
 - Patients who are taking medications for treatment of a disease for which the medication is not standard of care.



Evaluation

- Extremely useful 27
- Useful 61
- Somewhat useful 24
- Neutral 10
- Not useful 21



Lock-In

 Profiles Reviewed 	152
Total Cases	53
Case Rate	34.9%
Warning Cases	47
• Lock-In (LI) Cases	4
 Cases Continued in LI 	2
 Cases Removed from LI 	0



Questions?

