



STATE OF WEST VIRGINIA  
DEPARTMENT OF HEALTH AND HUMAN RESOURCES  
BUREAU FOR MEDICAL SERVICES



Office of Pharmacy Service  
Prior Authorization Criteria

**Austedo® (deutetrabenazine)**

**Effective 09/27/2018**

**[Prior Authorization Request Form](#)**

AUSTEDO is a vesicular monoamine transporter 2 (VMAT2) inhibitor indicated for the treatment of adults with chorea associated with Huntington's disease and for the treatment of tardive dyskinesia in adults.

**Initial\* Prior Authorization Criteria:**

- The patient must be within the age range as recommended by the FDA label; **AND**
- Patient must have been evaluated and found not to be suicidal or have untreated/undertreated depression; **AND**
- Patient must not be taking an MAOI (at least 14-days post-therapy), reserpine (must be >20 days post therapy) or any other concurrent VMAT2 inhibitor
- Prescriber must provide a brief description of the medical necessity of therapy by documenting all target symptoms and their impact on the patient's function and activities of daily living; **AND**

The following indication-specific criteria also apply:

**I. Treatment of Chorea associated with Huntington's Disease:**

1. Request must come from the treating neurologist; **AND**
2. All previous therapies must be documented along with their relative benefit. Unless contraindicated, the patient must have a documented 90-day trial of **amantadine** or **Xenazine (tetrabenazine)**.

**II. Treatment of Tardive Dyskinesia (TD):**

1. Request must come from the treating neurologist or psychiatrist; **AND**
2. Patient must provide a documented clinical diagnosis of tardive dyskinesia meeting DSM-V criteria including:
  - a. Involuntary athetoid or choreiform movements
  - b. History of treatment with a dopamine receptor blocking agent (DRBA) such as an antipsychotic or metoclopramide



STATE OF WEST VIRGINIA  
DEPARTMENT OF HEALTH AND HUMAN RESOURCES  
BUREAU FOR MEDICAL SERVICES



- c. Symptom duration lasting at least 8 weeks

**AND**

3. Prescriber must submit the results of an Abnormal Involuntary Movement Scale (AIMS) exam with every request for prior authorization of Austedo; **AND**
4. Prescriber must submit documentation of all other therapies attempted and their associated benefit (**including relevant AIMS scores**). Unless contraindicated, these therapies must include a **90-day trial with at least one** of the following agents: **clonazepam, amantadine, or Xenazine** (tetrabenazine).

**\*Initial prior-authorization will be for 90 days.  
Continuation of coverage requires clinically significant improvement in symptoms as compared to that seen using previous therapy.**

**References**

- 1.) Lexi-Comp drug monograph for Austedo (Reviewed 9/11/2018)
- 2.) Package insert for Austedo (last update 8/2017)
- 3.) Package insert for Xenazine (last update 9/2017)
- 4.) Abnormal Involuntary Movement Scale (AIMS) and Extrapyrimal Symptom Rating Scale (ESRS): cross-scale comparison in assessing tardive dyskinesia. Schizophr Res. 2005 Sep 15;77(2-3):119-28. Gharabawi GM<sup>1</sup>, Bossie CA, Lasser RA, Turkoz I, Rodriguez S, Chouinard G.
- 5.) UpToDate Tardive Dyskinesia: Prevention and Treatment. Article last updated July 24, 2017
- 6.) American Academy of Neurology Evidence-based guideline: Treatment of tardive syndromes. July 29, 2013.
- 7.) Treatment of Huntington's Disease. Neurotherapeutics. 2014 Jan; 11(1): 153–160.
- 8.) American Academy of Neurology Evidence-based guideline: Pharmacologic treatment of chorea in Huntington disease. August 7, 2012.