

EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ACNE AGENTS, TO	PICAL		
	ANTI-IN	FECTIVE	
	AKNE-MYCIN (erythromycin) AZELEX (azelaic acid) clindamycin erythromycin sodium sulfacetamide	ACZONE (dapsone) CLEOCIN-T (clindamycin) EVOCLIN (clindamycin) KLARON (sodium sulfacetamide)	Thirty (30) day trials each of one preferred retinoid and two unique chemical entities in two other subclasses, including the generic version of a requested non-preferred product, are required before a non- preferred agent will be authorized unless one of the exceptions on the PA form is present. (In cases of pregnancy, a trial of retinoids will not be required.)
	RETI	NOIDS	
	RETIN A liquid & Micro (tretinoin) TAZORAC (tazarotene) tretinoin cream, gel	AVITA DIFFERIN (adapalene) RETIN-A cream, gel (tretinoin)	PA required after 17 years of age for tretinoin products.
	KERATOLYTICS (Benzoyl Peroxides)	
	benzoyl peroxide ETHEXDERM (benzoyl peroxide) OSCION (benzoyl peroxide)	BENZAC WASH (benzoyl peroxide) BREVOXYL (benzoyl peroxide) DESQUAM (benzoyl peroxide) LAVOCLEN (benzoyl peroxide) TRIAZ (benzoyl peroxide)	Acne kits are non-preferred.
	COMBINAT	ION AGENTS	
	benzoyl peroxide/urea erythromycin/benzoyl peroxide sulfacetamide sodium/sulfur wash/cleanser	ACANYA (clindamycin phosphate/benzoyl peroxide) ^{NR} BENZACLIN GEL (benzoyl peroxide/clindamycin) BENZAMYCIN PAK (benzoyl peroxide/erythromycin) CLENIA (sulfacetamide sodium/sulfur)	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		DUAC CS (benzoyl peroxide/ clindamycin) INOVA 4/1 (benzoyl peroxide/salicylic acid) NUOX (benzoyl peroxide/sulfur) PLEXION (sulfacetamide sodium/sulfur) PRASCION (sulfacetamide sodium/sulfur) ROSAC (sulfacetamide sodium/avobenzone/sulfur) ROSADERM (sulfacetamide sodium/sulfur) ROSANIL (sulfacetamide sodium/sulfur) ROSULA (sulfacetamide sodium/sulfur) ROSULA (sulfacetamide sodium/sulfur) SULFOXYL (benzoyl peroxide/sulfur) SULFATOL (sulfacetamide sodium/sulfur/urea) ZIANA (clindamycin/tretinoin)	
ALZHEIMER'S AGE			
	CHOLINESTER	ASE INHIBITORS	
	ARICEPT (donepezil) ARICEPT ODT(donepezil) EXELON (rivastigmine)	COGNEX (tacrine) galantamine galantamine ER RAZADYNE (galantamine) RAZADYNE ER (galantamine)	A thirty (30) day trial of a preferred agent is required before a non- preferred agent In this class will be authorized unless one of the exceptions on the PA form is present.
	NMDA RECEPTO	DR ANTAGONIST	
	NAMENDA (memantine)		

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE 07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ANALGESICS, NAF	RCOTIC -SHORT ACTING (Non-p	arenteral)	
	APAP/codeine ASA/codeine codeine dihydrocodoeine/ APAP/caffeine hydrocodone/APAP hydrocodone/ibuprofen hydromorphone levorphanol morphine oxycodone oxycodone/APAP oxycodone/APAP pentazocine/APAP pentazocine/naloxone propoxyphene/APAP ROXICET (oxycodone/acetaminophen) tramadol tramadol/APAP VOPAC (codeine/acetaminophen)	ACTIQ (fentanyl) butalbital/APAP/caffeine/codeine butorphanol COMBUNOX (oxycodone/ibuprofen) DARVOCET (propoxyphene/APAP) DARVON (propoxyphene) DEMEROL (meperidine) DILAUDID (hydromorphone) fentanyl FENTORA (fentanyl) FIORICET W/ CODEINE (butalbital/APAP/caffeine/codeine) FIORINAL W/ CODEINE (butalbital/ASA/caffeine/codeine) LORCET (hydrocodone/APAP) LORTAB (hydrocodone/APAP) LORTAB (hydrocodone/APAP) LORTAB (hydrocodone/APAP) LORTAB (oxycodone/APAP) Meperidine OPANA (oxymorphone) oxycodone/ibuprofen OXYFAST (oxycodone) OXYIR (oxycodone) PANLOR (dihydrocodeine/ APAP/caffeine) PERCOCET (oxycodone/APAP) PERCODAN (oxycodone/APAP) PERCODAN (oxycodone/APAP) TALACEN (pentazocine/APAP) TALACEN (pentazocine/APAP) TALWIN NX (pentazocine/naloxone) TYLENOL W/CODEINE (APAP/codeine) ULTRACET (tramadol/APAP) VICOPROFEN (hydrocodone/APAP) VICOPROFEN (hydrocodone/APAP) ZYDONE (hydrocodone/APAP) ZYDONE (hydrocodone/APAP)	 Six (6) day trials of at least four (4) chemically distinct preferred agents (based on narcotic ingredient only), including the generic formulation of a requested non-preferred product, are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present. Fentanyl lozenges will only be approved for a diagnosis of cancer and as an adjunct to a long-acting agent. Fentanyl lozenges will not be approved for monotherapy. Limits: Unless the patient has escalating cancer pain or another diagnosis supporting increased quantities of short-acting opioids, all short acting solid forms of the narcotic analgesics are limited to 120 tablets per 30 days for the purpose of maximizing the use of longer acting medications to prevent unnecessary breakthrough pain in chronic pain therapy.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		XOLOX (oxycodone/APAP)	
ANALGESICS, NAF	RCOTIC - LONG ACTING (Non-pare	enteral)	
	DURAGESIC (fentanyl) KADIAN (morphine) methadone morphine ER OPANA ER (oxymorphone)	AVINZA (morphine) fentanyl MS CONTIN (morphine) ORAMORPH SR (morphine) oxycodone ER OXYCONTIN (oxycodone) ULTRAM ER (tramadol)	Six (6) day trials each of a total of four (4) preferred narcotic analgesics, including at least one long-acting agent, are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present. The generic form of the requested non-preferred agent, if available, must be tried before the non-preferred agent will be approved. Exception: Oxycodone ER will be authorized if a diagnosis of cancer is submitted without a trial of the preferred agents.
ANALGESICS, TOP	PICAL		
	capsaicin lidocaine lidocaine/prilocaine xylocaine	EMLA (lidocaine/prilocaine) FLECTOR PATCH (diclofenac) LIDODERM PATCH (lidocaine) LIDAMANTLE (lidocaine) LIDAMANTLE HC (lidocaine/hydrocortisone) LMX 4 (lidocaine) SYNERA (lidocaine/tetracaine) VOLTAREN GEL (diclofenac) ZOSTRIX (capsaicin)	DRAFT CRITERIA PENDING DRUG UTILIZATION COMMITTEE REVIEW ON 4/1/09. Ten (10) day trials of each of the preferred topical anesthetics (lidocaine, lidocaine/prilocaine, and xylocaine) are required before a non-preferred topical anesthetic will be approved unless one of the exceptions on the PA form is present. Lidoderm patches will be approved for a diagnosis of post-herpetic neuralgia.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE 07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
			Thirty (30) day trials of each of the preferred oral NSAIDS and capsaicin are required before Voltaren Gel will be approved unless one of the exceptions on the PA form is present.
			Flector patches will be approved only for a diagnosis of acute strain, sprain or injury after a five (5) day trial of one of the preferred oral NSAIDs and for a maximum duration of 14 days unless one of the exceptions on the PA forms is present.
ANDROGENIC AGI	ENTS		
	ANDRODERM (testosterone) ANDROGEL (testosterone)	TESTIM (testosterone)	The non-preferred agents will be approved only if one of the exceptions on the PA form is present.
ANGIOTENSIN MO	DULATORS		
	ACE INH	IBITORS	
	benazepril captopril enalapril fosinopril lisinopril quinapril ramipril	ACCUPRIL (quinapril) ACEON (perindopril) ALTACE (ramipril) CAPOTEN (captopril) LOTENSIN (benazepril) MAVIK (trandolapril) moexipril MONOPRIL (fosinopril) PRINIVIL (lisinopril) trandolapril UNIVASC (moexipril) VASOTEC (enalapril) ZESTRIL (lisinopril)	Fourteen (14) day trials of each of the preferred agents in the corresponding group, with the exception of the Direct Renin Inhibitors, are required before a non- preferred agent will be authorized unless one of the exceptions on the PA form is present.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
	ACE INHIBITOR COMBINATION DRUGS			
	benazepril/amlodipine benazepril/HCTZ captopril/HCTZ enalapril/HCTZ fosinopril/HCTZ lisinopril/HCTZ quinapril/HCTZ	ACCURETIC (quinapril/HCTZ) CAPOZIDE (captopril/HCTZ) LEXXEL (enalapril/felodipine) LOTENSIN HCT (benazepril/HCTZ) LOTREL (benazepril/amlodipine) moexipril/HCTZ MONOPRIL HCT (fosinopril/HCTZ) PRINZIDE (lisinopril/HCTZ) TARKA (trandolapril/verapamil) UNIRETIC (moexipril/HCTZ) VASERETIC (enalapril/HCTZ) ZESTORETIC (lisinopril/HCTZ)		
	ANGIOTENSIN II RECEP	TOR BLOCKERS (ARBs)		
	AVAPRO (irbesartan) BENICAR (olmesartan) COZAAR (losartan) 25mg DIOVAN (valsartan) MICARDIS (telmisartan)	ATACAND (candesartan) COZAAR (losartan) 50, 100mg TEVETEN (eprosartan)		
	ARB COM	BINATIONS		
	AVALIDE (irbesartan/HCTZ) AZOR (olmesartan/amlodipine) BENICAR-HCT (olmesartan/HCTZ) DIOVAN-HCT (valsartan/HCTZ) EXFORGE (valsartan/amlodipine) HYZAAR (losartan/HCTZ) MICARDIS-HCT (telmisartan/HCTZ)	ATACAND-HCT (candesartan/HCTZ) TEVETEN-HCT (erosartan/HCTZ)		
	DIRECT RENIN INHIBITORS			
	TEKTURNA (aliskiren) TEKTURNA HCT (aliskiren/HCTZ)		A thirty (30) day trial of one of a preferred ACE, ARB, or combination agents, at the maximum tolerable dose, is required before Tekturna will be approved.	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified. PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited

to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE 07/01/09

Version 2009.5

THERAPEUTIC	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
DRUG CLASS			
ANTICOAGULANTS	S, INJECTABLE ^{CL}		
	ARIXTRA (fondaparinux) FRAGMIN (dalteparin) LOVENOX (enoxaparin)	INNOHEP (tinzaparin)	Trials of each of the preferred agents will be required before a non- preferred agent will be approved unless one of the exceptions on the PA form is present.
ANTICONVULSAN	ſS		
	ADJU		
	carbamazepine CARBATROL (carbamazepine) DEPAKOTE ER (divalproex) DEPAKOTE SPRINKLE (divalproex) divalproex EC FELBATOL (felbamate) gabapentin GABITRIL (tiagabine) levetiracetam lamotrigine LYRICA (pregabalin) oxcarbazepine TOPAMAX (topiramate) valproic acid zonisamide	BANZEL(rufinamide) ^{NR} DEPAKENE (valproic acid) DEPAKOTE (divalproex) divalproex ER EPITOL (carbamazepine) EQUETRO (carbamazepine) KEPPRA XR (levetiracetam) LAMICTAL (lamotrigine) KEPPRA (levetiracetam) NEURONTIN (gabapentin) STAVZOR (valproic acid) TEGRETOL (carbamazepine) TEGRETOL XR (carbamazepine) TRILEPTAL (oxcarbazepine) VIMPAT (lacosamide) ^{NR} ZONEGRAN (zonisamide)	A fourteen (14) day trial of one of the preferred agents in the corresponding group is required for treatment naïve patients with a diagnosis of a seizure disorder before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present. A thirty (30) day trial of one of the preferred agents in the corresponding group is required for patients with a diagnosis other than seizure disorders unless one of the exceptions on the PA form is present. Keppra XR will be approved with a diagnosis of a seizure disorder with no trials of preferred agents required.
	BARBIT		
	mephobarbital phenobarbital primidone	MEBARAL (mephobarbital) MYSOLINE (primidone)	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
	BENZODI	AZEPINES		
	clonazepam DIASTAT (diazepam rectal) diazepam	KLONOPIN (clonazepam)		
	HYDAN	ITOINS		
	DILANTIN INFATABS (phenytoin) PEGANONE (ethotoin) phenytoin	CEREBYX (fosphenytoin) DILANTIN (phenytoin) PHENYTEK (phenytoin)		
	SUCCIN			
	CELONTIN (methsuximide) ethosuximide ZARONTIN (ethosuximide)			
ANTIDEPRESSANT	S, OTHER (second generation, no	n-SSRI)		
	bupropion SR bupropion XL CYMBALTA (duloxetine) EFFEXOR XR (venlafaxine) mirtazapine trazodone	APLENZIN (bupropion hbr) ^{NR} bupropion IR DESYREL (trazodone) EFFEXOR (venlafaxine) EMSAM (selegiline) nefazodone PRISTIQ (desvenlafaxine) REMERON (mirtazapine) venlafaxine venlafaxine ER WELLBUTRIN (bupropion) WELLBUTRIN SR (bupropion) WELLBUTRIN XL (bupropion)	A six (6) week trial each of a preferred agent and an SSRI is required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.	
ANTIDEPRESSANTS, SSRIs				
	citalopram fluoxetine fluvoxamine paroxetine sertraline	CELEXA (citalopram) LEXAPRO (escitalopram) LUVOX (fluvoxamine) LUVOX CR (fluvoxamine) PAXIL (paroxetine) PAXIL CR (paroxetine) paroxetine ER	Thirty (30) day trials each of two (2) of the preferred agents are required before a non-preferred agent will be approved unless one of the exceptions on the PA form is present. Upon hospital discharge, patients admitted with a primary	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		PEXEVA (paroxetine) PROZAC (fluoxetine) RAPIFLUX (fluoxetine) SARAFEM (fluoxetine) ZOLOFT (sertraline)	mental health diagnosis and have been stabilized on a non-preferred SSRI will receive an authorization to continue that drug.
ANTIEMETICS			
	5HT3 RECEPT	OR BLOCKERS	
	ondansetron ondansetron ODT	ANZEMET (dolasetron) KYTRIL (granisetron) granisetron SANCUSO (granisetron) ZOFRAN (ondansetron) ZOFRAN ODT (ondansetron)	A 3-day trial of a preferred agent is required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present. PA is required for all agents when limits are exceeded.
	CANNA	BINOIDS	
		CESAMET (nabilone) MARINOL (dronabinol)	Cesamet will be authorized only for the treatment of nausea and vomiting associated with cancer chemotherapy for patients who have failed to respond adequately to 3- day trials of conventional treatments such as promethazine or ondansetron and are over 18 years of age. Marinol will be authorized only for the treatment of anorexia associated with weight loss in patients with AIDS or cancer and unresponsive to megestrol, the prophylaxis of chemotherapy induced nausea and vomiting unresponsive to 3-day trials of ondansetron or promethazine and for patients between the ages of 18 and 65 years of age.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified. PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited

to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		ANTAGONISTS	
	EMEND (aprepitant)		
ANTIFUNGALS, OF	RAL		
	clotrimazole fluconazole* ketoconazole ^{CL} nystatin terbinafine ^{CL}	ANCOBON (flucytosine) DIFLUCAN (fluconazole) griseofulvin GRIS-PEG (griseofulvin) itraconazole LAMISIL (terbinafine) MYCELEX (clotrimazole) MYCOSTATIN Tablets (nystatin) NIZORAL (ketoconazole) NOXAFIL (posaconazole) SPORANOX (itraconazole) VFEND (voriconazole)	Non-preferred agents will be approved only if one of the exceptions on the PA form is present. *PA is required when limits are exceeded. PA is not required for griseofulvin suspension for children up to 6 years of age for the treatment of tinea capitis.
ANTIFUNGALS, TO			
	ANTIFU econazole MENTAX (butenafine) NAFTIN (naftifine) nystatin	ingals ciclopirox ERTACZO (sertaconazole) EXELDERM (sulconazole) LOPROX (ciclopirox) MYCOSTATIN (nystatin) NIZORAL (ketoconazole) OXISTAT (oxiconazole) PENLAC (ciclopirox) SPECTAZOLE (econazole) VUSION (miconazole/petrolatum/zinc oxide) XOLEGEL (ketoconazole)	Fourteen (14) day trials of two (2) of the preferred agents are required before one of the non-preferred agents will be authorized unless one of the exceptions on the PA form is present. If a non-preferred shampoo is requested, a fourteen (14) day trial of one preferred product (ketoconazole shampoo) is required.) Oxistat cream will be approved for children 12 and under for tinea corporis, tinea cruris, tinea pedis, and tinea (pityriasis) versicolor.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	ANTIFUNGAL/STER	OID COMBINATIONS	
	clotrimazole/betamethasone nystatin/triamcinolone	LOTRISONE (clotrimazole/betamethasone) MYCOLOG (nystatin/triamcinolone)	
ANTIHISTAMINES,	MINIMALLY SEDATING		
	ANTIHIS	TAMINES	
	ALAVERT (loratadine) cetirizine (OTC) loratadine TAVIST-ND (loratadine)	ALLEGRA (fexofenadine) CLARINEX Tablets (desloratadine) CLARINEX REDITABS (desloratadine) CLARINEX Syrup (desloratadine) CLARITIN (loratadine) fexofenadine XYZAL (levocetirizine) ZYRTEC (Rx and OTC) (cetirizine) ZYRTEC SYRUP (Rx and OTC) (cetirizine)	Thirty (30) day trials of at least two (2) chemically distinct preferred agents (in the age appropriate form), including the generic formulation of a requested non-preferred product, are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.
	ANTIHISTAMINE/DECONO	SESTANT COMBINATIONS	
	ALAVERT-D (loratadine/pseudoephedrine) cetirizine/pseudoephedrine (OTC) loratadine/pseudoephedrine SEMPREX-D (acrivastine/ pseudoephedrine)	ALLEGRA-D (fexofenadine/pseudoephedrine) CLARINEX-D (desloratadine/pseudoephedrine) CLARITIN-D (loratadine/pseudoephedrine) ZYRTEC-D (Rx and OTC) (cetirizine/pseudoephedrine)	
ANTIMIGRAINE AG	ENTS, TRIPTANS		
	TRIP	TANS	
	IMITREX (sumatriptan) MAXALT MLT (rizatriptan) RELPAX (eletriptan)	AMERGE (naratriptan) AXERT (almotriptan) FROVA (frovatriptan) MAXALT (rizatriptan) sumatriptan ZOMIG (zolmitriptan)	Three (3) day trials each of the preferred agents are required before a non-preferred agent will be approved unless one of the exceptions on the PA form is present. Quantity limits apply for this drug class.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	TREXIMET (sumatriptan/naproxen sodium)		
ANTIPARKINSON'S	S AGENTS (Oral)		
	ANTICHOL	INERGICS	
	benztropine KEMADRIN (procyclidine) trihexyphenidyl	COGENTIN (benztropine)	Patients starting therapy on drugs in this class must show a documented allergy to all of the preferred agents, in the corresponding class, before a non-preferred agent will be authorized.
	COMT IN	HIBITORS	
		COMTAN (entacapone) TASMAR (tolcapone)	
	DOPAMINE	AGONISTS	
	ropinirole	MIRAPEX (pramipexole) REQUIP (ropinirole) REQUIP XL (ropinirole)	Mirapex, Requip, and Requip XL will be approved for a diagnosis of Parkinsonism with no trials of preferred agents required.
	OTHER ANTIPARK	(INSON'S AGENTS	
	amantadine bromocriptine carbidopa/levodopa selegiline STALEVO (levodopa/carbidopa/entacapone)	AZILECT (rasagiline) ELDEPRYL (selegiline) levodopa/carbidopa ODT PARCOPA (levodopa/carbidopa) SINEMET (levodopa/carbidopa) ZELAPAR (selegiline)	
ANTIPSYCHOTICS	, ATYPICAL (Oral)		
	clozapine GEODON (ziprasidone) INVEGA (paliperidone) RISPERDAL SOLUTION (risperidone)	ABILIFY (aripiprazole) CLOZARIL (clozapine) FAZACLO (clozapine) RISPERDAL (risperidone)	A fourteen (14) day trial of a preferred agent is required for treatment naïve patients before a non-preferred agent will be approved

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE 07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	risperidone SEROQUEL (quetiapine) SEROQUEL XR (quetiapine)	risperidone solution ZYPREXA (olanzapine)	unless one of the exceptions on the PA form is present. Upon discharge, a hospitalized patient stabilized on a non-preferred agent may receive authorization to continue this drug for labeled indications and at recommended dosages.
			Abilify will be prior authorized for MDD if the following criteria are met:
			 The patient is at least 18 year of age. Diagnosis of Major Depressive Disorder (MDD) not responsive to other antidepressants. Evidence of trials of appropriate therapeutic duration at a maximum tolerable dose of at least two (2) of the following agents: Selective Serotonin Reuptake Inhibitors (SSRI), Norepinephrine Reuptake Inhibitors, or bupropion. Prescribed in conjunction with an SSRI, SNRI or bupropion. The daily dose does not exceed 15 mg.
	ATYPICAL ANTIPSYCHO	TIC/SSRI COMBINATIONS	
		SYMBYAX (olanzapine/fluoxetine)	
ANTIVIRALS (Oral)			
	ANTI H	ERPES	
	acyclovir VALTREX (valacyclovir)	famciclovir FAMVIR (famciclovir) ZOVIRAX (acyclovir)	Five (5) day trials each of the preferred agents are required before the non-preferred agents will be authorized unless one of the

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified. PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited

to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
			exceptions on the PA form is present.
	ANTI INF	LUENZA	
		FLUMADINE (rimantadine) RELENZA (zanamivir) rimantadine SYMMETREL (amantadine) TAMIFLU (oseltamivir)	The anti influenza agents will be approved only for a diagnosis of influenza.
ATOPIC DERMATIT	ris		
	ELIDEL (pimecrolimus) PROTOPIC (tacrolimus)		
BETA BLOCKERS	(Oral)		
		OCKERS	
	acebutolol atenolol betaxolol bisoprolol metoprolol ER nadolol propranolol propranolol ER sotalol timolol TOPROL XL (metoprolol)	BETAPACE (sotalol) BLOCADREN (timolol) BYSTOLIC (nebivolol) CARTROL (carteolol) CORGARD (nadolol) INDERAL LA (propranolol) INNOPRAN XL (propranolol) KERLONE (betaxolol) LEVATOL (penbutolol) LOPRESSOR (metoprolol) SECTRAL (acebutolol) TENORMIN (atenolol) ZEBETA (bisoprolol)	Fourteen (14) day trials each of three (3) chemically distinct preferred agents, including the generic formulation of a requested non-preferred product, are required before one of the non-preferred agents will be approved unless one of the exceptions on the PA form is present.
		IC COMBINATION DRUGS	
	atenolol/chlorthalidone bisoprolol/HCTZ metoprolol/HCTZ nadolol/bendroflumethiazide propranolol/HCTZ	CORZIDE (nadolol/bendroflumethiazide) INDERIDE (propranolol/HCTZ) LOPRESSOR HCT (metoprolol/HCTZ) TENORETIC (atenolol/chlorthalidone) ZIAC (bisoprolol/HCTZ)	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
	BETA- AND ALF	PHA-BLOCKERS		
	carvedilol labetalol	COREG (carvedilol) COREG CR (carvedilol) TRANDATE (labetalol)		
BLADDER RELAX	ANT PREPARATIONS			
	DETROL LA (tolterodine) ENABLEX (darifenacin) oxybutynin oxybutynin ER SANCTURA (trospium) SANCTURA XR (trospium) VESICARE (solifenacin)	DETROL (tolterodine) DITROPAN (oxybutynin) DITROPAN XL (oxybutynin) OXYTROL (oxybutynin)	A thirty (30) day trial each of the chemically distinct preferred agents is required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.	
BONE RESORPTIO	N SUPPRESSION AND RELATED	AGENTS		
	BISPHOSF	PHONATES		
	alendronate FOSAMAX PLUS D (alendronate/vitamin D)	ACTONEL (risedronate) ACTONEL WITH CALCIUM (risedronate/calcium) BONIVA (ibandronate) DIDRONEL (etidronate) FOSAMAX (alendronate)	A 30-day trial of one of the preferred agents is required before a non- preferred agent will be authorized unless one of the exceptions on the PA form is present.	
	OTHER BONE RESORPTION SUPP	RESSION AND RELATED AGENTS		
	MIACALCIN (calcitonin)	calcitonin EVISTA (raloxifene) FORTEO (teriparatide) FORTICAL (calcitonin)	Evista will be approved for postmenopausal women with osteoporosis or at high risk for invasive breast cancer.	
BPH AGENTS				
	AVODART (dutasteride) finasteride	PROSCAR (finasteride) RAPAFLO (silodosin) ^{NR}	Thirty (30) day trials each of at least two (2) chemically distinct preferred agents, including the generic formulation of a requested non- preferred agent, are required before	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
			a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.
	ALPHA B	LOCKERS	
	doxazosin FLOMAX (tamsulosin) terazosin UROXATRAL (alfuzosin)	CARDURA (doxazosin) CARDURA XL (doxazosin) HYTRIN (terazosin) RAPAFLO (silodosin) ^{NR}	
BRONCHODILATO	RS, ANTICHOLINERGIC		
	ANTICHO	LINERGIC	
	ATROVENT HFA (ipratropium) ipratropium SPIRIVA (tiotropium)		Thirty (30) day trials each of the preferred agents in the corresponding group are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.
	ANTICHOLINERGIC-BETA	AGONIST COMBINATIONS	
	COMBIVENT (albuterol/ipratropium)	albuterol/ipratropium DUONEB (albuterol/ipratropium)	For severely compromised patients, albuterol/ipratropium will be approved if the combined volume of albuterol and ipratropium nebules is inhibitory.
BRONCHODILATO	RS, BETA AGONIST		
		N SOLUTION	
	albuterol	ACCUNEB (albuterol)** BROVANA (arformoterol) metaproterenol PERFOROMIST (formoterol) PROVENTIL (albuterol)	Thirty (30) day trials each of the chemically distinct preferred agents in their corresponding groups are required before a non-preferred agent in that group will be authorized

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		XOPENEX (levalbuterol)	unless one of the exceptions on the PA form is present.
			**No PA is required for ACCUNEB for children up to 5 years of age.
	INHALERS, L	ONG-ACTING	
	FORADIL (formoterol) SEREVENT (salmeterol)		
	INHALERS, SI	HORT-ACTING	
	MAXAIR (pirbuterol) PROAIR HFA (albuterol) PROVENTIL HFA (albuterol) VENTOLIN HFA (albuterol)	ALUPENT (metaproterenol) PROVENTIL (albuterol) XOPENEX HFA (levalbuterol)	Xopenex Inhalation Solution will be approved for 12 months for a diagnosis of asthma or COPD for patients on concurrent asthma controller therapy (either oral or inhaled) with documentation of failure on a trial of albuterol or documented intolerance of albuterol, or for concurrent diagnosis of heart disease.
	OF	AL	
	albuterol terbutaline	BRETHINE (terbutaline) metaproterenol VOSPIRE ER (albuterol)	
CALCIUM CHANNE	EL BLOCKERS		
	LONG-	ACTING	
	amlodipine diltiazem felodipine ER nifedipine ER nisoldipine verapamil ER	ADALAT CC (nifedipine) CALAN SR (verapamil) CARDENE SR (nicardipine) CARDIZEM CD, LA, SR (diltiazem) COVERA-HS (verapamil) DILACOR XR (diltiazem) DYNACIRC CR (isradipine) ISOPTIN SR (verapamil) NORVASC (amlodipine) PLENDIL (felodipine)	Fourteen (14) day trials each of the preferred agents are required before a non-preferred agent will be approved unless one of the exceptions on the PA form is present.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified. PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited

to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		PROCARDIA XL (nifedipine) SULAR (nisoldipine) TIAZAC (diltiazem) VERELAN/VERELAN PM (verapamil)	
	SHORT	ACTING	
	diltiazem verapamil	ADALAT (nifedipine) CALAN (verapamil) CARDENE (nicardipine) CARDIZEM (diltiazem) DYNACIRC (isradipine) isradipine nicardipine nimodipine nifedipine NIMOTOP (nimodipine) PROCARDIA (nifedipine)	
CEPHALOSPORIN	S AND RELATED ANTIBIOTICS (O	-	
	BETA LACTAMS AND BETA LACTAM/BET	A-LACTAMASE INHIBITOR COMBINATIONS	
	amoxicillin/clavulanate AUGMENTIN XR (amoxicillin/clavulanate)	MOXATAG (amoxicillin) ^{NR}	Five (5) day trials each of the preferred agents required before a non-preferred agent is authorized unless one of the exceptions on the PA form is present.
	CEPHALC	DSPORINS	
	cefaclor cefadroxil cefdinir cefpodoxime cefprozil cefuroxime cephalexin SPECTRACEF (cefditoren)	CECLOR (cefaclor) CEDAX (ceftibuten) CEFTIN (cefuroxime) CEFZIL (cefprozil) DURICEF (cefadroxil) KEFLEX (cephalexin) OMNICEF (cefdinir) PANIXINE (cephalexin) RANICLOR (cefaclor) SUPRAX (cefixime) VANTIN (cefpodoxime)	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA		
DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGEN 15			
	^t GENERATION ANTIHISTAMINES				
		, 1 ST GENERATION			
	chlorpheniramine maleate	brompheniramine maleate			
	clemastine	brompheniramine tannate			
	cyproheptadine	BROVEX (brompheniramine tannate)			
	diphenhydramine	carbinoxamine maleate			
	promethazine	LODRANE (brompheniramine maleate and			
		tannate)			
		LOHIST (brompheniramine maleate)			
		PALGIC (carbinoxamine maleate)			
		TANACOF (brompheniramine tannate)			
		TANAHIST-PD (chorpheniramine tannate)			
	ANTITUSSIVE-ANTIHIS	AMINE COMBINATIONS			
	codeine/promethazine				
	dextromethorphan HBR/promethazine				
	ANTIHISTAMINE-ANTITUSSIVE-D	ECONGESTANT COMBINATIONS			
	brompheniramine/dextromethorphan				
	HBR/pseudoephedrine				
	chlorpheniramine/dextromethorphan/				
	pseudoephedrine				
	promethazine/codeine/phenylephrine				
	ANTITUSSIVE-DECONG	ESTANT COMBINATIONS			
		MUCINEX-D (guaifenesin/pseudoephedrine)			
	DECONGESTANTS				
	phenylephrine	NASOP (phenylephrine)			
	pseudoephedrine				

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	ANTITUSSIVES/EXPECTORANTS		
	<mark>benzonatate</mark> guaifenesin guaifenesin/dextromethorphan	MUCINEX (guaifenesin) MUCINEX-DM (guaifenesin/dextromethorphan) TESSALON (benzonatate)	
	DECONGESTANT-ANTIHISTAMINE-	ANTICHOLINERGIC COMBINATIONS	
	phenylephrine/chlorpheniramine/ scopolamine	DURAHIST (pseudoephedrine/chlorpheniramine/ methscopolamine) EXTENDRYL CHW /JR TAB (phenylephrine/chlorpheniramine/ scopolamine) EXTENDRYL SOL (phenylephrine/dexchlorpheniramine/ methscopolamine) NOHIST-PLUS (phenylephrine/ chlorpheniramine/methscopolamine) phenylephrine/chlorpheniramine/ methscopolamine pseudoephedrine/chlorpheniramine/ methscopolamine phenylephrine/dexchlorpheniramine/ methscopolamine RE-DRYLEX JR (phenylephrine/ chlorpheniramine/scopolamine) RE-DRYLEX SYRUP (phenylephrine/dexchlorpheniramine/ methscopolamine) SCOPOHIST (pseudoephedrine/ chlorpheniramine/methscopolamine)	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA		
	DECONGESTANT-ANTIHISTAMINE COMBINATIONS				
		STAMINE COMBINATIONS BROVEX-D (phenylephrine/ brompheniramine) CHLOR-TAN SUSP (phenylephrine tannate/pyrilamine tannate/ chlorpheniramine) DURATUSS DA (pseudoephedrine/chlorpheniramine) DYTAN-D CHW/SUSP (phenylephrine tannate/diphenhydramine tannate) LODRANE 12D/24D//D (pseudoephedrine/brompheniramine) LOHIST 12D/PD (pseudoephedrine/brompheniramine) LOHIST 12D/PD (pseudoephedrine/chlorpheniramine) NALEX-A LIQUID/SUSPENSION (phenylephrine/phenyltoloxamine/ chlorpheniramine) phenylephrine/brompheniramine phenylephrine/torompheniramine phenylephrine/brompheniramine	PA CRITERIA		
		pyrilamine/chlorpheniramine) RONDEC (phenylephrine/chlorpheniramine) RU-HIST FORTE (phenylephrine/pyrilamine/ chlorpheniramine)			
		RYNATAN (phenylephrine/chlorpheniramine) SUDAL 12 (pseudoephedrine/chlorpheniramine) TANNATE PED SUSP (phenylephrine/chlorpheniramine)			

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
CYTOKINE & CAM	ANTAGONISTS CL		
	CIMZIA (certolizumab/pegol) ENBREL (etanercept) HUMIRA (adalimumab) KINERET (anakinra) RAPTIVA (efalizumab)		
ERYTHROPOIESIS	S STIMULATING PROTEINS CL		
	PROCRIT (rHuEPO)	ARANESP (darbepoetin) EPOGEN (rHuEPO)	A thirty (30) day trial of the preferred agent is required before a non- preferred agent will be authorized unless one of the exceptions on the PA form is present.
FLUOROQUINOLO	NES, ORAL		
	AVELOX (moxifloxacin) CIPRO (ciprofloxacin) Suspension ciprofloxacin ciprofloxacin ER LEVAQUIN (levofloxacin)	CIPRO (ciprofloxacin) Tablets CIPRO XR (ciprofloxacin) FACTIVE (gemifloxacin) FLOXIN (ofloxacin) NOROXIN (norfloxacin) ofloxacin PROQUIN XR (ciprofloxacin)	A five (5) day trial of one of the preferred agents is required before a non-preferred agent will be approved unless one of the exceptions on the PA form is present.
GENITAL WARTS	AGENTS		
	ALDARA (imiquimod)	CONDYLOX (podofilox) podofilox VEREGEN (sinecatechins)	A thirty (30) day trial of the preferred agent is required before a non- preferred agent will be authorized unless one of the exceptions on the PA form is present.
GLUCOCORTICOI	DS, INHALED		
	GLUCOCO	DRTICOIDS	
	AEROBID (flunisolide) AEROBID-M (flunisolide) ASMANEX (mometasone) AZMACORT (triamcinolone)	ALVESCO (ciclesonide) budesonide PULMICORT (budesonide)	Thirty (30) day trials each of the preferred agents are required before a non-preferred agent will be authorized unless one of the

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
	FLOVENT HFA (fluticasone) FLOVENT Diskus (fluticasone) QVAR (beclomethasone)		exceptions on the PA form is present. Pulmicort Respules do not require a prior authorization for children through 8 years of age or for individuals unable to use an MDI. When children who have been stabilized on Pulmicort Respules reach age 9, prescriptions for the Pulmicort inhaler will be authorized for them.	
	GLUCOCORTICOID/BRONCH	ODILATOR COMBINATIONS		
	ADVAIR (fluticasone/salmeterol) ADVAIR HFA (fluticasone/salmeterol) SYMBICORT(budesonide/formoterol)			
GROWTH HORMON	NE ^{CL}			
	GENOTROPIN (somatropin) NUTROPIN (somatropin) NUTROPIN AQ (somatropin)	HUMATROPE (somatropin) INCRELEX (mecasermin) NORDITROPIN (somatropin) OMNITROPE (somatropin) SAIZEN (somatropin) SEROSTIM (somatropin) TEV-TROPIN (somatropin) ZORBTIVE (somatropin)	The preferred agents must be tried before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present. Patients already on a non-preferred agent will receive authorization to continue therapy on that agent for the duration of the existing PA.	
HEPATITIS B TREATMENTS				
	EPIVIR HBV (lamivudine) HEPSERA (adefovir) TYZEKA (telbivudine)	BARACLUDE (entecavir)	A thirty (30) day trial of one of the preferred agents is required before the non-preferred agent will be authorized unless one of the exceptions on the PA form is present.	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE 07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
HEPATITIS C TREA	ATMENTS ^{CL}		
	PEGASYS (pegylated interferon) PEG-INTRON (pegylated interferon) ribavirin	COPEGUS (ribavirin) INFERGEN (consensus interferon) REBETOL (ribavirin)	Patients starting therapy in this class must try the preferred agent of a dosage form before a non-preferred agent of that dosage form will be authorized.
HYPOGLYCEMICS	, INCRETIN MIMETICS/ENHANCER	S	
	BYETTA (exenatide) JANUMET (sitagliptin/metformin) JANUVIA (sitagliptin) SYMLIN (amylin)		Byetta and Symlin are both subject to the following step therapy edits: Byetta-Current history of therapy with a sulfonylurea, thiazolidinedione (TZD), and/or metformin. No gaps of therapy greater than 30 days in the past 180 days. Symlin- History of insulin utilization in the past 90 days. No gaps in therapy of greater than 30 days.
HYPOGLYCEMICS	, INSULINS		
	HUMALOG (insulin lispro) vials only HUMALOG MIX (insulin lispro/lispro protamine) vials only HUMULIN (insulin) vials only LANTUS (insulin glargine) all forms LEVEMIR (insulin detemir) all forms NOVOLIN (insulin) all forms NOVOLOG (insulin aspart) all forms NOVOLOG MIX all forms (insulin aspart/aspart protamine)	APIDRA (insulin glulisine) HUMALOG PEN/KWIKPEN (insulin lispro) HUMALOG MIX PENS (insulin lispro/lispro protamine) HUMULIN PEN (insulin)	 To receive Apidra, patients must meet the following criteria: be 4 years or older; be currently on a regimen including a longer-acting or basal insulin. have had a trial of a similar preferred agent, Novolog or Humalog, with documentation that the desired results were not achieved. Current prescriptions for Humalog Pens and cartridges, Humalog Kwikpens, Humalog Mix Pens, and

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified. PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited

to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
			Humulin Pens will be grandfathered.	
HYPOGLYCEMICS	, MEGLITINIDES			
	STARLIX (nateglinide)	PRANDIN (repaglinide)	A thirty (30) day triial of the preferred agent is required before a non- preferred agent will be authorized, unless one of the exceptions on the PA form is present.	
HYPOGLYCEMICS	, TZDS			
		INEDIONES		
	ACTOS (pioglitazone) AVANDIA (rosiglitazone)			
		BINATIONS		
	ACTOPLUS MET (pioglitazone/metformin) AVANDAMET (rosiglitazone/metformin) AVANDARYL (rosiglitazone/glimepiride) DUETACT (pioglitazone/glimepiride)			
IMPETIGO AGENTS	S, TOPICAL			
	ALTABAX (retapamulin) mupirocin bacitracin gentamicin sulfate	BACTROBAN (mupirocin) CORTISPORIN (bacitracin/neomycin/polymyxin/HC)	Ten (10) day trials of at least one preferred agent, including the generic formulation of a requested non-preferred agent, are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.	
INTRANASAL RHINITIS AGENTS				
	ANTICHOL	INERGICS		
		ATROVENT(ipratropium) ipratropium	Thirty (30) day trials of one preferred agent in the antihistamine and corticosteroid groups are required	
	the listing of a particular brand or generic name inclu agents apply in addition to general Drug Utilization I			

to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
			before an anti-cholinergic agent will be approved unless one of the exceptions on the PA form is present.
	ANTIHIS	TAMINES	
	ASTELIN (azelastine) PATANASE (olopatadine)	ASTEPRO (azelastine)	DRAFT CRITERIA PENDING DRUG UTILIZATION COMMITTEE REVIEW ON 4/1/09. Thirty (30) day trials of both preferred intranasal antihistamines and a thirty (30) day trial of one of the preferred intranasal corticosteroids are required before the non-preferred agent will be approved unless one of the exceptions on the PA form is present.
	CORTICO	STEROIDS	
	fluticasone propionate NASACORT AQ (triamcinolone) NASONEX (mometasone) VERAMYST (fluticasone furoate)	BECONASE AQ (beclomethasone) flunisolide FLONASE (fluticasone propionate) NASALIDE (flunisolide) NASAREL (flunisolide) RHINOCORT AQUA (budesonide)	Thirty (30) day trials of each preferred agent in the corticosteroid group are required before a non- preferred corticosteroid agent will be authorized unless one of the exceptions on the PA form is present.
LEUKOTRIENE MO			
	ACCOLATE (zafirlukast) SINGULAIR (montelukast)	ZYFLO (zileuton)	Thirty (30) day trials each of the preferred agents are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
LIPOTROPICS, OTI	HER (non-statins)		
	BILE ACID SE	QUESTRANTS	
	cholestyramine colestipol	COLESTID (colestipol) QUESTRAN (cholestyramine) WELCHOL (colesevelam)	A twelve (12) week trial of one of the preferred agents is required before a non-preferred agent in the corresponding category will be authorized.
			Zetia, as monotherapy, will only be approved for patients who cannot take statins or other preferred agents.
			Welchol will be approved for add-on therapy only after an insufficient response to the maximum tolerable dose of a statin after 12 weeks of therapy.
	CHOLESTEROL ABS	ORPTION INHIBITORS	
		ZETIA (ezetimibe)	Zetia will be approved for add-on therapy only after an insufficient response to the maximum tolerable dose of a statin after 12 weeks of therapy.
	FATTY	ACIDS	
	LOVAZA (omega-3-acid ethyl esters)		DRAFT CRITERIA UNTIL APPROVED BY DUR Lovaza will be approved for the treatment of high triglyceride levels (> 400mg/dL) when the patient is intolerant or not responsive to, or not a candidate for nicotinic acid or fibrate therapy.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified. PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited

to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	FIBRIC ACID DERIVATIVES		
	fenofibrate gemfibrozil TRICOR (fenofibrate) <mark>TRILIPIX (fenofibrate)^{NR}</mark>	ANTARA (fenofibrate) FENOGLIDE (fenofibrate) LOFIBRA (fenofibrate) LOPID (gemfibrozil) TRIGLIDE (fenofibrate)	
	NIA	CIN	
	niacin NIASPAN (niacin)	NIACELS (niacin) NIADELAY (niacin) SLO-NIACIN (niacin)	
LIPOTROPICS, ST	ATINS		
,,		TINS	
	CRESTOR (rosuvastatin) LESCOL (fluvastatin) LESCOL XL (fluvastatin) LIPITOR (atorvastatin) lovastatin pravastatin simvastatin	ALTOPREV (lovastatin) MEVACOR (lovastatin) PRAVACHOL (pravastatin) ZOCOR (simvastatin)	Twelve (12) week trials each of two (2) of the preferred statins, including the generic formulation of a requested non-preferred agent, are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.
	STATIN CON	MBINATIONS	
	ADVICOR (lovastatin/niacin) CADUET (atorvastatin/amlodipine) SIMCOR (simvastatin/niacin ER)	VYTORIN (simvastatin/ ezetimibe)	Vytorin will be approved only after an insufficient response to the maximum tolerable dose of Lipitor (atorvastatin) or Crestor (rosuvastatin) after 12 weeks, unless one of the exceptions on the PA form is present.
MACROLIDES/KET	OLIDES (Oral)		•
	KETEK (telithromycin)		Requests for telithromycin will be authorized if there is documentation of the use of any antibiotic within the

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
			past 28 days.
	MACD	OLIDES	
			Five (E) day trials each of the
	azithromycin clarithromycin erythromycin	BIAXIN (clarithromycin) BIAXIN XL (clarithromycin) clarithromycin ER E.E.S. (erythromycin ethylsuccinate) E-MYCIN (erythromycin) ERYC (erythromycin) ERYPED (erythromycin ethylsuccinate) ERY-TAB (erythromycin) ERYTHROCIN (erythromycin stearate) erythromycin estolate PCE (erythromycin) ZITHROMAX (azithromycin) ZMAX (azithromycin)	Five (5) day trials each of the preferred agents are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.
MULTIPLE SCLER	OSIS AGENTS CL		
	AVONEX (interferon beta-1a) BETASERON (interferon beta-1b) COPAXONE (glatiramer) REBIF (interferon beta-1a)	TYSABRI (natalizumab)	A 30-day trial of a preferred agent will be required before a non- preferred agent will be approved. Tysabri will only be approved for members who are enrolled in the TOUCH Prescribing Program.
MUSCLE RELAXA	NTS, ORAL		
	ACUTE MUSCULOSKELE	TAL RELAXANT AGENTS	
	chlorzoxazone cyclobenzaprine methocarbamol	AMRIX (cyclobenzaprine) carisoprodol carisoprodol/ASA carisoprodol/ASA/codeine FEXMID (cyclobenzaprine) FLEXERIL (cyclobenzaprine) methocarbamol/ASA orphenadrine orphenadrine/ASA/caffeine	DRAFT CRITERIA PENDING DRUG UTILIZATION COMMITTEE REVIEW ON 4/1/09. Thirty (30) day trials of the preferred acute musculoskeletal relaxants are required before a non-preferred acute musculoskeletal agent will be approved, with the exception of

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE 07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		PARAFON FORTE DSC (chlorzoxazone) ROBAXIN (methocarbamol) SKELAXIN (metaxalone) SOMA (carisoprodol) SOMA COMPOUND (carisoprodol /ASA) SOMA COMP w/ COD (carisoprodol/ASA/codeine)	carisoprodol. Thirty (30) day trials of the preferred acute musculoskeletal relaxants and Skelaxin are required before carisoprodol will be approved.
	MUSCULOSKELETAL RELAXANT	AGENTS USED FOR SPASTICITY	
	baclofen dantrolene tizanidine	DANTRIUM (dantrolene) ZANAFLEX (tizanidine)	Thirty (30) day trials of the preferred skeletal muscle relaxants associated with the treatment of spasticity (are required before non-preferred agents will be approved unless one of the exceptions on the PA form is present.
NSAIDS			
	NONSEI	ECTIVE	
	diclofenac etodolac fenoprofen flurbiprofen ibuprofen (Rx and OTC) INDOCIN (indomethacin) (suspension only) indomethacin ketorolac naproxen (Rx only) oxaprozin piroxicam sulindac	ADVIL (ibuprofen) ANAPROX (naproxen) ANSAID (flurbiprofen) CATAFLAM (diclofenac) CLINORIL (sulindac) DAYPRO (oxaprozin) FELDENE (piroxicam) INDOCIN (indomethacin) ketoprofen LODINE (etodolac) meclofenamate mefenamic acid MOTRIN (ibuprofen) nabumetone NALFON (fenoprofen) NAPRELAN (naproxen) NAPROSYN (naproxen) NUPRIN (ibuprofen) ORUDIS (ketoprofen)	Thirty (30) day trials of each of the preferred agents are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		PONSTEL (meclofenamate) tolmetin VOLTAREN (diclofenac)	
	NSAID/GI PROTECT	ANT COMBINATIONS	
		ARTHROTEC (diclofenac/misoprostol) PREVACID/NAPRAPAC (naproxen/lansoprazole)	
	COX-II SE	ELECTIVE	
	CELEBREX (celecoxib) ^{CL} meloxicam	MOBIC (meloxicam)	Celebrex will be approved for patients with a GI Risk Score of ≥13.
OPHTHALMIC ANT	TBIOTICS		
	ciprofloxacin ofloxacin VIGAMOX (moxifloxacin)	AZASITE (azithromycin) CILOXAN (ciprofloxacin) OCUFLOX (ofloxacin) QUIXIN (levofloxacin) ZYMAR (gatifloxacin)	Five (5) day trials each of the preferred agents are required before non-preferred agents will be authorized unless one of the exceptions on the PA form is present.
OPTHALMIC ANTI-	INFLAMMATORIES		
	ACULAR/LS/PF (ketorolac) flurbiprofen NEVANAC (nepafenac) XIBROM (bromfenac)	diclofenac DUREZOL (difluprednate)	DRAFT CRITERIA PENDING DRUG UTILIZATION COMMITTEE REVIEW ON 4/1/09. Five (5) day trials of each of the preferred ophthalmic anti- inflammatory agents are required before nonpreferred agens will be authorized unless one of the exceptions on the PA form is present.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified. PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
DRUG CLASS	R ALLERGIC CONJUNCTIVITIS		
	ACULAR (ketorolac) ALAWAY (ketotifen) ALREX (loteprednol) cromolyn OPTIVAR (azelastine) PATADAY (olopatadine) PATANOL (olopatadine) ZADITOR OTC (ketotifen)	ALAMAST (pemirolast) ALOCRIL (nedocromil) ALOMIDE (lodoxamide) CROLOM (cromolyn) ELESTAT (epinastine) EMADINE (emedastine) ketotifen OPTICROM (cromolyn)	Thirty (30) day trials each of two (2) of the preferred agents are required before non-preferred agents will be authorized, unless one of the exceptions on the PA form is present.
OPHTHALMICS, GL	AUCOMA AGENTS		
	COMBINATI	ON AGENTS	
	COSOPT (dorzolamide/timolol)	COMBIGAN (brimonidine/timolol) dorzolamide/timolol	Authorization for a non-preferred agent will only be given if there is an allergy to the preferred agents.
	BETA BL	OCKERS	
	Betaxolol BETOPTIC S (betaxolol) carteolol levobunolol metipranolol timolol	BETAGAN (levobunolol) BETIMOL (timolol) ISTALOL (timolol) OPTIPRANOLOL (metipranolol) TIMOPTIC (timolol)	
	CARBONIC ANHY	DRASE INHIBITORS	
	AZOPT (brinzolamide) TRUSOPT (dorzolamide)	dorzolamide	
	PARASYMPA	THOMIMETICS	
	CARBOPTIC (carbachol) ISOPTO CARBACHOL (carbachol) PHOSPHOLINE IODIDE (echothiophate iodide) Pilocarpine	ISOPTO CARPINE (pilocarpine) PILOPINE HS (pilocarpine)	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	LUMIGAN (bimatoprost) TRAVATAN (travoprost) TRAVATAN-Z (travoprost)	XALATAN (latanoprost)	
	SYMPATHO	OMIMETICS	
	ALPHAGAN P (brimonidine) brimonidine dipivefrin	ALPHAGAN (brimonidine) PROPINE (dipivefrin)	
OTIC FLUOROQUII	NOLONES		
	CIPRODEX (ciprofloxacin/dexamethasone) ofloxacin	CIPRO HC (ciprofloxacin/hydrocortisone) FLOXIN (ofloxacin)	Five (5) day trials each of the preferred agents are required before a non-preferred agent will be approved unless one of the exceptions on the PA form is present.
PANCREATIC ENZ	YMES		
	CREON PANCRECARB ULTRASE ULTRASE MT VIOKASE	KUZYME LIPRAM PALCAPS PANCREASE PANGESTYME PANOKASE PLARETASE	Thirty (30) day trials each of at least three (3) preferred agents are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present. Non-preferred agents will be approved for members with cystic fibrosis.
PARATHYROID AG			
	ergocalciferol calcitriol HECTOROL (doxercalciferol) ZEMPLAR (paricalcitol)	DRISDOL (ergocalciferol) ROCALTROL (calcitriol) SENSIPAR (cinacalcet)	A thirty (30) day trial of a preferred agent will be required before a non- preferred agent will be approved.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA		
PEDICULICIDES/SO	PEDICULICIDES/SCABICIDES, TOPICAL				
	EURAX (crotamiton) OVIDE (malathion) permethrin (Rx and OTC) pyrethrins-piperonyl butoxide	lindane	Trials of the preferred agents (which are age and weight appropriate) are required before lindane will be approved unless one of the exceptions on the PA form is present.		
PHOSPHATE BIND	ERS				
	FOSRENOL (lanthanum) PHOSLO (calcium acetate) RENAGEL (sevelamer)	calcium acetate ELIPHOS (calcium acetate) RENVELA (sevelamer carbonate)	Thirty (30) day trials of at least two (2) preferred agents are required unless one of the exceptions on the PA form is present.		
PLATELET AGGRE	EGATION INHIBITORS				
	AGGRENOX (dipyridamole/ASA) cilostazol PLAVIX (clopidogrel)	dipyridamole PERSANTINE (dipyridamole) PLETAL (cilostazol) TICLID (ticlopidine) ticlopidine	A thirty (30) day trial of a preferred agent is required before a non- preferred agent will be approved unless one of the exceptions on the PA form is present.		
PRENATAL VITAM	INS				
	prenatal vitamin 27 w/calcium/ferrous fumarate/folic acid prenatal vitamins 28 w/calcium/iron ps complex/folic acid prenatal vitamins/ferrous fumarate/docusate/folic acid prenatal vitamins/ferrous fumarate/folic acid prenatal vitamins/ferrous fumarate/folic acid/selenium prenatal vitamins/iron, carbonyl/folic acid prenatal vitamins/iron polysaccharides complex/folic acid prenatal vitamins comb 10/ferrous fumarate/folic acid prenatal vitamin no. 15/iron, carbonyl/folic	CARENATAL DHA CITRANATAL DHA COMBI RX FOLBECAL DUET/DUET DHA FOLTABS PLUS DHA NATACHEW NATAFORT NATELLE PLUS W/DHA NEEVO NOVANATAL OB-NATAL ONE OPTINATE PRECARE/PRECARE PREMIER PREMESIS			

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	acid/docusate sod prenatal vitamin no. 16/iron, carbonyl/folic acid/docusate sod prenatal vitamin no. 17/iron, carbonyl/folic acid/docusate sod prenatal vitamin w-o calcium/ferrous fumarate/folic acid prenatal vitamins w-o vit A/iron, carbonyl/folic acid	prenatal vitamins/ferrous bis-glycinate chelate/folic acid prenatal vitamins/iron, carbonyl/omega- 3/FA/fat combo no. 1 prenatal vitamins comb no. 20/iron bisgly/folic acid/DHA prenatal vitamins no. 22/iron, carbonyl/FA/docusate/DHA prenatal vitamins w-CA, FE, FA (<1 mg) prenatal vitamins w-CA, FE, FA (<1 mg) prenatal vitamins w-o calcium/iron ps complex/FA prenatal vitamins w-o CA no. 5/ferrous fumarate/folic acid prenatal vitamins w-o CA no. 2 prenatal vitamins w-o calcium no. 9/iron/folic acid PRENATE DHA/PRENATE ELITE PRENAVITE PRENAVITE PRENAZA PRIMACARE RENATE/RENATE DHA SELECT-OB TANDEM DHA/TANDEM OB	
PROTON PUMP IN			
	NEXIUM (esomeprazole) PREVACID Capsules (lansoprazole)	ACIPHEX (rabeprazole) KAPIDEX (dexlansoprazole) NEXIUM PACKETS (esomeprazole) omeprazole pantoprazole PREVACID Solu-Tabs (lansoprazole) PREVACID Suspension (lansoprazole) PRILOSEC (omeprazole) PROTONIX (pantoprazole) ZEGERID (omeprazole/sodium bicarbonate)	Thirty (30) day trials each of the preferred agents are required before a non-preferred agent will be approved unless one of the exceptions on the PA form is present. Prior authorization is not required for Prevacid Solu-Tabs for patients ≤8 years of age.

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified. PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited

to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
PULMONARY ANTI	HYPERTENSIVES-ENDOTHELIN F		
	TRACLEER (bosentan)	LETAIRIS (ambrisentan)	DRAFT CRITERIA PENDING DRUG UTILIZATION COMMITTEE REVIEW ON 4/1/09. These agents will only be approved for the treatment of pulmonary artery hypertension World Health Organization (WHO) group I. Letairis will only be approved for patients with WHO class II or III symptoms after a fourteen (14) day trial of the preferred agent unless one of the exceptions on the PA form is present. Users of Letairis as of 3/31/09 will be allowed to continue therapy with that drug.
SEDATIVE HYPNO	TICS		
	BENZODI	AZEPINES	
	temazepam	DALMANE (flurazepam) DORAL (quazepam) estazolam flurazepam HALCION (triazolam) PROSOM (estazolam) RESTORIL (temazepam) triazolam	Fourteen (14) day trials of the preferred agents in both categories are required before a non-preferred agent will be authorized unless one of the exceptions on the PA form is present.
	zolpidem	AMBIEN (zolpidem) AMBIEN CR (zolpidem) AQUA CHLORAL (chloral hydrate) chloral hydrate	

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE 07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA		
		LUNESTA (eszopiclone) ROZEREM (ramelteon) SOMNOTE (chloral hydrate) SONATA (zaleplon) zaleplon			
STIMULANTS AND RELATED AGENTS					
	4				
	ADDERALL XR (amphetamine salt combination) amphetamine salt combination dextroamphetamine VYVANSE (lisdexamphetamine)	ADDERALL (amphetamine salt combination) DESOXYN (methamphetamine) DEXEDRINE (dextroamphetamine) DEXTROSTAT (dextroamphetamine)	 Except for Strattera, PA is required for adults >18 years. One of the preferred agents in each group (amphetamines and non-amphetamines) must be tried for thirty (30) days before a non-preferred agent will be authorized. Thirty (30) day trials of at least three (3) antidepressants are required before amphetamines will be approved for depression. Provigil will only be approved for patients >16 years of age with a diagnosis of narcolepsy. Strattera will not be approved for concurrent administration with amphetamines or methylphenidates, except for 30 days or less for tapering purposes. Strattera is limited to a maximum of 100mg per day. 		
NON-AMPHETAMINE					
	CONCERTA (methylphenidate) FOCALIN (dexmethylphenidate) FOCALIN XR (dexmethylphenidate) METADATE CD (methylphenidate)	DAYTRANA (methylphenidate) dexmethylphenidate METADATE ER (methylphenidate) pemoline			

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.



EFFECTIVE

07/01/09

Version 2009.5

THERAPEUTIC DRUG CLASS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA		
	methylphenidate methylphenidate ER STRATTERA (atomoxetine)	PROVIGIL (modafinil) RITALIN (methylphenidate) RITALIN LA (methylphenidate) RITALIN-SR (methylphenidate)			
ULCERATIVE COLITIS AGENTS					
	OR				
	ASACOL (mesalamine) COLAZAL (balsalazide) DIPENTUM (olsalazine) LIALDA (mesalamine) PENTASA (mesalamine) sulfasalazine	APRISO (mesalamine) [№] AZULFIDINE (sulfasalazine) balsalazide	Thirty (30) day trials of each of the preferred agents of a dosage form must be tried before a non-preferred agent of that dosage form will be authorized unless one of the exceptions on the PA form is present.		
RECTAL					
	CANASA (mesalamine) mesalamine	ROWASA (mesalamine)			
MISC BRAND/GENERIC					
	SANDOSTATIN (octreotide)	octreotide	Thirty (30) day trials each of the preferred agents are required before a non-preferred agent will be authorized.		

Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC are not covered unless specified.

PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.

^{CL} – Requires Clinical PA. For detailed clinical criteria, please refer to: <u>http://www.wvdhhr.org/bms/sPharmacy/Drugs/bms_drugs_main.asp</u>.