

EFFECTIVE 07/01/2018 Version 2018.3e

- Prior authorization for a non-preferred agent in any class will be given only if there has been a trial of the preferred brand/generic equivalent or preferred formulation of the active ingredient, at a therapeutic dose, that resulted in a partial response with a documented intolerance.
- Prior authorization of a non-preferred isomer, pro-drug, or metabolite will be considered with a trial of a preferred parent drug of the same chemical entity, at a therapeutic dose, that resulted in a partial response with documented intolerance or a previous trial and therapy failure, at a therapeutic dose, with a preferred drug of a different chemical entity indicated to treat the submitted diagnosis. (The required trial may be overridden when documented evidence is provided that the use of these preferred agent(s) would be medically contraindicated.)
- Unless otherwise specified, the listing of a particular brand or generic name includes all legend forms of that drug. OTC drugs are not covered unless specified.
- PA criteria for non-preferred agents apply in addition to general Drug Utilization Review policy that is in effect for the entire pharmacy program, including, but not limited to, appropriate dosing, duplication of therapy, etc.
- The use of pharmaceutical samples will not be considered when evaluating the members' medical condition or prior prescription history for drugs that require prior authorization.
- Quantity limits may apply. Refer to the Limits List on the BMS Website by clicking the hyperlink.
- Unless otherwise indicated, non-preferred combination products require medical reasoning beyond convenience or enhanced compliance as to why the clinical need cannot be met with a preferred agent or combination of preferred single-ingredient agents.
- Acronyms
  - CL Requires clinical PA. For detailed clinical criteria, please go to the PA criteria page by clicking the hyperlink.
  - NR Denotes a new drug which has not yet been reviewed by the P & T Committee. These agents are available only on appeal to the BMS Medical Director.
  - o AP Non-preferred and selected preferred drugs, where indicated, are subject to auto-PA criteria. See PA criteria column.



EFFECTIVE 07/01/2018 Version 2018.3e

CLASSES CHANGING	Status Changes	PA Criteria Changes	New Drugs
ACNE AGENTS, TOPICAL – ROSACEA AGENTS		XXXX	XXXX
ANTIDEPRESSANTS, SSRI's			XXXX
ANTIHYPUREMICS, URICOSURIC-XOI COMBINATION	XXXX		XXXX
ANTIRETROVIRALS, PROTEASE INHIBITORS			XXXX
ANTIRETROVIRALS, INSTI & NRTI COMBINATION			XXXX
COPD AGENTS, ANTICHOLINERGIC/BETA-AGONIST/GLUCOCORTICOID COMBINATION		XXXX	XXXX
FLUOROQUINOLONES, ORAL			XXXX
GLUCOCORTICOIDS, INHALED			XXXX
HYPOGLYCEMICS, INSULIN AND RELATED AGENTS			XXXX
IMMUNOMODULATORS, ATOPIC DERMATITIS		XXXX	
IRRITABLE BOWEL SYNDROME, SHORT BOWEL SYNDROME AND SELECTED GI AGENTS			XXXX
PLATELET AGGREGATION INHIBITORS	XXXX		XXXX
STIMULANTS, NON-AMPHETAMINE	XXXX	XXXX	XXXX



EFFECTIVE 07/01/2018 Version 2018.3e

This is not an all-inclusive list of available covered drugs and includes only managed categories. Refer to cover page for complete list of rules governing this PDL.

THERAPEUTIC DRUG CLASS

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
ACNE AGENTS, TOPICALAP			
CLASS PA CRITERIA: Non-preferred agents require a thirty (30) day trial of one (1) preferred retinoid and two (2) unique chemical entites in two (2) other subclasses, including the generic version of the requested non-preferred product, before they will be approved, unless one (1) of the exceptions on the PA form is present.			
In cases of pregnancy, a trial of retinoids will <i>not</i> be Acne kits are non-preferred.	e required. For members eighteen (18) years of age	e or older, a trial of retinoids will not be required.	
Specific Criteria for sub-class will be listed bel day trial of all preferred agents in that sub-class.	ow. NOTE: Non-preferred agents in the Rosacea su	ub-class are available only on appeal and require at least a 30-	
au, mai or an protonou agonto in marcazo ciasa.	ANTI-INFECTIVE		
clindamycin gel, lotion, medicated swab, solution erythromycin gel, solution	ACZONE (dapsone) AKNE-MYCIN (erythromycin) AZELEX (azelaic acid) CLEOCIN-T (clindamycin) CLINDACIN PAC (clindamycin) CLINDAGEL (clindamycin) clindamycin foam erythromycin medicated swab EVOCLIN (clindamycin) FABIOR (tazarotene) KLARON (sulfacetamide) OVACE/PLUS (sulfacetamide) sodium sulfacetamide 10% cleansing gel sulfacetamide cleanser sulfacetamide cleanser ER sulfacetamide suspension		
	RETINOIDS		
RETIN-A (tretinoin) TAZORAC (tazarotene)	adapalene ATRALIN (tretinoin) AVITA (tretinoin) DIFFERIN (adapalene) RETIN-A MICRO (tretinoin) tretinoin cream, gel tretinoin gel micro	In addition to the Class Criteria: PA required for members eighteen (18) years of age or older.	
D 0.0T0 4004	KERATOLYTICS		
benzoyl peroxide cleanser Rx & OTC, 10% cream OTC, gel Rx & OTC, lotion OTC, wash OTC	BENZEFOAM ULTRA (benzoyl peroxide) BENZEPRO (benzoyl peroxide) benzoyl peroxide cloths, medicated pads, microspheres cleanser BP 10-1 (benzoyl peroxide)		



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	SS
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	BP WASH 7% LIQUID PACNEX/HP/LP (benzoyl peroxide) PANOXYL-4, -8 OTC (benzoyl peroxide) PERSA-GEL OTC (benzoyl peroxide) SULPHO-LAC (sulfur)	
	COMBINATION AGENTS	
erythromycin/benzoyl peroxide	ACANYA (clindamycin phosphate/benzoyl peroxide)  AVAR/-E/LS (sulfur/sulfacetamide)  BENZACLIN GEL (benzoyl peroxide/ clindamycin)  BENZAMYCIN PAK (benzoyl peroxide/ erythromycin)  benzoyl peroxide/clindamycin gel benzoyl peroxide/urea  CERISA (sulfacetamide sodium/sulfur)  CLARIFOAM EF (sulfacetamide/sulfur)  CLENIA (sulfacetamide sodium/sulfur)  DUAC (benzoyl peroxide/clindamycin)  EPIDUO (adapalene/benzoyl peroxide)*  INOVA 4/1, 5/2benzoyl peroxide/salicylic acid)  NEUAC (clindamycin phosphate/benzoyl peroxide)  NUOX (benzoyl peroxide/sulfur)  ONEXTON (clindamycin phosphate/benzoyl peroxide)  PRASCION (sulfacetamide sodium/sulfur)  SE 10-5 SS (sulfacetamide/sulfur)  SSS 10-4 (sulfacetamide /sulfur)  SSS 10-5 foam (sulfacetamide /sulfur)  sulfacetamide sodium/sulfur cloths, lotion, pads, suspension  sulfacetamide/sulfur wash/cleanser  sulfacetamide/sulfur wash kit  sulfacetamide/sulfur wash kit	In addition to the Class Criteria: Non-preferred combination agents require thirty (30) day trials of the corresponding preferred single agents before they will be approved.  *PA required for combination agents with Retinoid products for members eighteen (18) years of age or older.



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLA	ASS
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	ROSACEA AGENTS	
FINACEA GEL (azelaic acid) MIRVASO GEL (brimonidine) metronidazole cream metronidazole gel (NDCs 00115-1474-46, 00168-0275-45, 00713-0637-37, 51672- 4116-06, 66993-0962-45 only)	FINACEA FOAM (azelaic acid) METROCREAM (metronidazole) METROGEL GEL (metronidazole) METROLOTION (metronidazole) metronidazole lotion metronidazole gel (all other NDCs) NORITATE CREAM (metronidazole) RHOFADE (oxymetazoline) ROSADAN (metronidazole) SOOLANTRA CREAM (metronidazole)	Subclass criteria: Non-preferred agents are available only on appeal and require evidence of 30-day trials of all chemically-unique preferred agents in the sub-class.
ALZHEIMER'S AGENTSAP		
the exceptions on the PA form is present.	ts require a thirty (30) day trial of a preferred agent in to to forty-five (45) years of age if there is no diagnosis of	the same sub-class before they will be approved, unless one (1) of of Alzheimer's disease.
	CHOLINESTERASE INHIBITORS	s
donepezil 5 and 10 mg	ARICEPT (donepezil) donepezil 23 mg* EXELON CAPSULE (rivastigmine) EXELON PATCH (rivastigmine) galantamine galantamine ER RAZADYNE (galantamine) RAZADYNE ER (galantamine) rivastigmine	*Donepezil 23 mg tablets will be authorized if the following criteria are met:  1. There is a diagnosis of moderate-to-severe Alzheimer's Disease <b>and</b> 2. There has been a trial of donepezil 10 mg daily for at least three (3) months and donepezil 20 mg daily for an additional one (1) month.
	NMDA RECEPTOR ANTAGONIS	
memantine	NAMENDA (memantine) NAMENDA XR (memantine)*	*Namenda XR requires ninety (90) days of compliant therapy with Namenda.
СНО	LINESTERASE INHIBITOR/NMDÁ RECEPTOR ANTA	AGONIST COMBINATIONS
	NAMZARIC (donepezil/memantine)	Combination agents require thirty (30) day trials of each corresponding preferred single agent.
ANALGESICS, NARCOTIC LON	G ACTING (Non-parenteral) <sup>AP</sup>	
CLASS PA CRITERIA: Non-preferred agen requested non-preferred agent (if available) the requested non-preferred brand agent, the	ts require six (6) day trials of two (2) chemically distinct before they will be approved, unless one (1) of the exce en another generic non-preferred agent must be triale	et preferred agents <b>AND</b> a six (6) day trial of the generic form of the eptions on the PA form is present. If no generic form is available for ed instead. <b>NOTE: All long-acting opioid agents require a prior</b> and indication and specify previous opioid and non-opioid therapies
buprenorphine patch (labeler 00093 only) BUTRANS (buprenorphine) EMBEDA (morphine/naltrexone)	ARYMO ER (morphine sulfate) BELBUCA (buprenorphine buccal film)* buprenorphine patch (all labelers excl 00093)	*Belbuca prior authorization requires manual review. Full PA criteria may be found on the <u>PA Criteria</u> page by clicking the hyperlink.



EFFECTIVE 07/01/2018 Version 2018.3e

This is not an all-inclusive list of available covered drugs and includes only managed categories. Refer to cover page for complete list of rules governing this PDL.

THERAPEUTIC DRUG CLASS			
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
fentanyl transdermal 12, 25, 50, 75, 100 mcg/hr morphine ER tablets	CONZIP ER (tramadol) DOLOPHINE (methadone) DURAGESIC (fentanyl) EXALGO ER (hydromorphone) fentanyl transdermal 37.5, 62.5, 87.5 mcg/hr hydromorphone ER HYSINGLA ER (hydrocodone) KADIAN (morphine) LAZANDA SPRAY (fentanyl) methadone** MORPHABOND ER (morphine sulfate) morphine ER capsules (generic for Avinza) morphine ER capsules (generic for Kadian) MS CONTIN (morphine) NUCYNTA ER (tapentadol) OPANA ER (oxymorphone) oxycodone ER** OXYCONTIN (oxycodone) oxymorphone ER** tramadol ER*** ULTRAM ER (tramadol) XARTEMIS XR (oxycodone/ acetaminophen) XTAMPZA ER (oxycodone) ZOHYDRO ER (hydrocodone)	**Methadone, oxycodone ER and oxymorphone ER will be authorized without a trial of the preferred agents if a diagnosis of cancer is submitted.  ***Tramadol ER requires a manual review and may be authorized for ninety (90) days with submission of a detailed treatment plan including anticipated duration of treatment and scheduled follow-ups with the prescriber.	
ANIAL OFFICE NADCOTIC CHOPT	A OTINIO (NI IV-s		

### ANALGESICS, NARCOTIC SHORT ACTING (Non-parenteral)<sup>AP</sup>

CLASS PA CRITERIA: Non-preferred agents require six (6) day trials of at least four (4) chemically distinct preferred agents (based on the narcotic ingredient only), including the generic formulation of the requested non-preferred agent, before they will be approved, unless one (1) of the exceptions on the PA form is present.

NOTE: All tramadol and codeine products require a prior authorization for children under 18 years of age. Requests must be for an FDA approved age and indication and specify non-opioid therapies attempted.

indication and specify non-opioid therapies atten	npted.	
APAP/codeine	ABSTRAL (fentanyl)	Fentanyl buccal, nasal and sublingual products will only be
butalbital/APAP/caffeine/codeine	ACTIQ (fentanyl)	authorized for a diagnosis of cancer and as an adjunct to a
codeine	butalbital/ASA/caffeine/codeine	long-acting agent. These dosage forms will not be authorized
hydrocodone/APAP 2.5/325 mg, 5/325 mg,	butorphanol	for monotherapy.
7.5/325 mg,10/325 mg	CAPITAL W/CODEINE (APAP/codeine)	
hydrocodone/APAP solution	DEMEROL (meperidine)	Limits: Unless the patient has escalating cancer pain or
hydrocodone/ibuprofen	dihydrocodeine/ APAP/caffeine	another diagnosis supporting increased quantities of short-
hydromorphone tablets	DILAUDID (hydromorphone)	acting opioids, all short acting solid forms of the narcotic
morphine	fentanyl	analgesics are limited to 120 tablets per thirty (30) days.
oxycodone tablets, concentrate, solution	FENTORA (fentanyl)	Longer-acting medications should be maximized to prevent
oxycodone/APAP	FIORICET W/ CODEINE	unnecessary breakthrough pain in chronic pain therapy.
oxycodone/ASA	(butalbital/APAP/caffeine/codeine)	
tramadol	FIORINAL W/ CODEINE	Immediate-release tramadol is limited to 240 tablets per thirty



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
tramadol/APAP	(butalbital/ASA/caffeine/codeine) hydrocodone/APAP 5/300 mg, 7.5/300 mg, 10/300 mg hydromorphone liquid, suppositories IBUDONE (hydrocodone/ibuprofen) LAZANDA (fentanyl) levorphanol LORCET (hydrocodone/APAP) LORTAB (hydrocodone/APAP) meperidine NORCO (hydrocodone/APAP) NUCYNTA (tapentadol) ONSOLIS (fentanyl) OPANA (oxymorphone) OXECTA (oxycodone) oxycodone capsules oxycodone/ibuprofen oxymorphone pentazocine/naloxone PERCOCET (oxycodone/APAP) PRIMLEV (oxycodone/APAP) PRIMLEV (oxycodone/APAP) REPREXAIN (hydrocodone/ibuprofen) ROXICODONE (oxycodone) RYBIX ODT (tramadol) SUBSYS (fentanyl) SYNALGOS-DC (dihydrocodeine/ASA/ caffeine) TYLENOL W/CODEINE (APAP/codeine) ULTRACET (tramadol/APAP) ULTRAM (tramadol) VERDROCET (hydrocodone/APAP) VICODIN (hydrocodone/APAP) VICOPROFEN (hydrocodone/APAP) VICOPROFEN (hydrocodone/ibuprofen) XODOL (hydrocodone/ibuprofen) XYLON (hydrocodone/ibuprofen) ZAMICET (hydrocodone/APAP)	(30) days.
ANDROGENIC AGENTS  CLASS PA CRITERIA: A pop-preferred agent wi	Il only be authorized if one (1) of the exceptions on th	e PA form is present
ANDRODERM (testosterone)	ANDROID (methyltestosterone)	o i Atlanti la prodotti.
ANDROGEL (testosterone)	AVEED VIAL (testosterone undecanoate)	
METHITEST (methyltestosterone)	AXIRON (testosterone)	
testosterone cypionate vial <sup>CL</sup>	FORTESTA (testosterone)	
testosterone enanthate vial <sup>CL</sup>	methyltestosterone capsule	



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	SS
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	NATESTO (testosterone) STRIANT BUCCAL (testosterone) TESTIM (testosterone) TESTRED (methyltestosterone) testosterone gel VOGELXO (testosterone)	
ANESTHETICS, TOPICALAP		
<b>CLASS PA CRITERIA:</b> Non-preferred agents re- PA form is present.	quire ten (10) day trials of each preferred agent befo	ore they will be approved, unless one (1) of the exceptions on the
lidocaine lidocaine/prilocaine xylocaine	EMLA (lidocaine/prilocaine) LIDAMANTLE (lidocaine) LIDAMANTLE HC (lidocaine/hydrocortisone) lidocaine/hydrocortisone LIDOTRAL CREAM (lidocaine) SYNERA (lidocaine/tetracaine) VOPAC MDS (ketoprofen/lidocaine)	
ANGIOTENSIN MODULATORSAP		
CLASS PA CRITERIA: Non-preferred agents re Inhibitors, before they will be approved, unless on		ent in the same sub-class, with the exception of the Direct Renin
	ACE INHIBITORS	
benazepril captopril enalapril fosinopril lisinopril quinapril ramipril	ACCUPRIL (quinapril) ACEON (perindopril) ALTACE (ramipril) EPANED (enalapril)* LOTENSIN (benazepril) MAVIK (trandolapril) moexipril perindopril PRINIVIL (lisinopril) QBRELIS SOLUTION (lisinopril)** trandolapril UNIVASC (moexipril) VASOTEC (enalapril) ZESTRIL (lisinopril)	*Epaned will be authorized with a diagnosis of hypertension, symptomatic heart failure or asymptomatic left ventricular dysfunction provided that the patient is less than seven (7) years of age <b>OR</b> is unable to ingest a solid dosage form due to documented oral-motor difficulties or dysphagia.  **Qbrelis solution may be authorized for children ages 6-10 who are unable to tolerate a solid dosage form. Qbrelis may also be authorized for older patients with clinical documentation indicating oral-motor difficulties or dysphagia.
benazepril/amlodipine	ACE INHIBITOR COMBINATION DRUG ACCURETIC (quinapril/HCTZ)	<del>სა</del>
benazeprii/HCTZ captoprii/HCTZ enalaprii/HCTZ	CAPOZIDE (captopril/HCTZ) LOTENSIN HCT (benazepril/HCTZ) LOTREL (benazepril/amlodipine)	



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	SS
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
fosinopril/HCTZ lisinopril/HCTZ quinapril/HCTZ	moexipril/HCTZ PRESTALIA (perindopril/amlodipine) PRINZIDE (lisinopril/HCTZ) TARKA (trandolapril/verapamil) trandolapril/verapamil VASERETIC (enalapril/HCTZ)	
	ZESTORETIC (lisinopril/HCTZ)  ANGIOTENSIN II RECEPTOR BLOCKERS	(ARRs)
irbesartan losartan valsartan olmesartan	ATACAND (candesartan) AVAPRO (irbesartan) BENICAR (olmesartan) candesartan COZAAR (losartan) DIOVAN (valsartan) EDARBI (azilsartan) eprosartan MICARDIS (telmisartan) telmisartan TEVETEN (eprosartan)	(AILDS)
	ARB COMBINATIONS	
ENTRESTO (valsartan/sucubitril) <sup>CL*</sup> irbesartan/HCTZ losartan/HCTZ olmesartan/amlodipine olmesartan/amlodipine/HCTZ olmesartan/HCTZ valsartan/amlodipine valsartan/HCTZ	ATACAND-HCT (candesartan/HCTZ) AVALIDE (irbesartan/HCTZ) AZOR (olmesartan/amlodipine) BENICAR-HCT (olmesartan/HCTZ) BYVALSON (nebivolol/valsartan) candesartan/HCTZ DIOVAN-HCT (valsartan/HCTZ) EDARBYCLOR (azilsartan/chlorthalidone) EXFORGE (valsartan/amlodipine) EXFORGE HCT (valsartan/amlodipine/HCTZ) HYZAAR (losartan/HCTZ) MICARDIS-HCT (telmisartan/HCTZ) telmisartan/amlodipine telmisartan HCTZ TEVETEN-HCT (eprosartan/HCTZ) TRIBENZOR (olmesartan/amlodipine/HCTZ) TWYNSTA (telmisartan/amlodipine) valsartan/amlodipine/HCTZ	*Entresto will only be authorized for patients diagnosed with chronic heart-failure (NYHA classification 2-4) with an EF ≤ 40%.
	DIRECT RENIN INHIBITORS	
	AMTURNIDE (aliskiren/amlodipine/HCTZ) TEKAMLO (aliskiren/amlodipine) TEKTURNA (aliskiren) TEKTURNA HCT (aliskiren/HCTZ)	Substitute for Class Criteria: Tekturna requires a thirty (30) day trial of one (1) preferred ACE, ARB, or combination agent, at the maximum tolerable dose, before it will be authorized unless one (1) of the exceptions on the PA form is present.



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS			
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
	VALTURNA (aliskiren/valsartan)	Amturnide, Tekamlo, Tekturna HCT or Valturna will be authorized if the criteria for Tekturna are met and the patient also needs the other agents in the combination.	
ANTIANGINAL & ANTI-ISCHEMIC			
or a combination agent containing one (1) of these		sium channel blocker, a beta blocker, or a nitrite as single agents	
RANEXA (ranolazine) <sup>AP</sup>			
ANTIBIOTICS, GI & RELATED AGE	NTS		
CLASS PA CRITERIA: Non-preferred agents red PA form is present.	uire a fourteen (14) day trial of a preferred agent bef	ore they will be approved, unless one (1) of the exceptions on the	
metronidazole tablet neomycin tinidazole	ALINIA (nitazoxanide) DIFICID (fidaxomicin)* FLAGYL (metronidazole) FLAGYL ER (metronidazole ER) metronidazole capsule paromomycin TINDAMAX (tinidazole) VANCOCIN (vancomycin) vancomycin** XIFAXAN (rifaximin)***	*Dificid will be authorized if the following criteria are met:  1. There is a diagnosis of severe <i>C. difficile</i> infection; and  2. There is no response to prior treatment with vancomycin for ten (10) to fourteen (14) days.  **Vancomycin will be authorized for treatment of mild to moderate <i>C. difficile</i> infections after a fourteen (14) day trial of metronidazole. Severe <i>C. difficile</i> infections do not require a trial of metronidazole for authorization.  ***Full PA criteria may be found on the PA Criteria page by clicking the hyperlink.	
ANTIBIOTICS, INHALED		onothing the ripportant.	
•		and documentation of therapeutic failure before they will be	
BETHKIS (tobramycin) KITABIS PAK (tobramycin)	CAYSTON (aztreonam) TOBI (tobramycin) TOBI PODHALER (tobramycin) tobramycin		
ANTIBIOTICS, TOPICAL	•		
CLASS PA CRITERIA: Non-preferred agents require ten (10) day trials of at least one preferred agent, including the generic formulation of the requested non-preferred agent, before they will be approved, unless one (1) of the exceptions on the PA form is present.			
bacitracin (Rx, OTC) gentamicin sulfate mupirocin ointment	BACTROBAN (mupirocin) CENTANY (mupirocin) CORTISPORIN (bacitracin/neomycin/polymyxin/HC) mupirocin cream neomycin/polymyxin/pramoxine		



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ANTIBIOTICS, VAGINAL		
approved, unless one (1) of the exceptions on the	PA form is present.	at the manufacturer's recommended duration, before they will be
clindamycin cream CLINDESSE (clindamycin) metronidazole	AVC (sulfanilamide) CLEOCIN CREAM (clindamycin) CLEOCIN OVULE (clindamycin) METROGEL (metronidazole) NUVESSA (metronidazole) VANDAZOLE (metronidazole)	
ANTICOAGULANTS		
CLASS PA CRITERIA: Non-preferred agents req	uire a trial of each preferred agent in the same sub-	class, unless one (1) of the exceptions on the PA form is present.
	INJECTABLECL	
enoxaparin	ARIXTRA (fondaparinux) fondaparinux FRAGMIN (dalteparin) LOVENOX (enoxaparin)	
	ORAL	
COUMADIN (warfarin) ELIQUIS (apixaban) PRADAXA (dabigatran) warfarin XARELTO (rivaroxaban)	SAVAYSA (edoxaban)	
ANTICONVULSANTS		
	e disorder, non-preferred agents require a fourteen (1 tions on the PA form is present; patients currently on	14) day trial of a preferred agent in the same sub-class before established therapies shall be grandfathered.
For all other diagnoses, non-preferred agents requexceptions on the PA form is present.	uire a thirty (30) day trial of a preferred agent in the sa	ame sub-class before they will be approved, unless one (1) of the
In situations where AB-rated generic equivalent problems brand name product to be reimbursed.	oducts are available, "Brand Medically Necessary" m	nust be hand-written by the prescriber on the prescription for the
	ADJUVANTS	
carbamazepine carbamazepine ER carbamazepine XR	APTIOM (eslicarbazepine) BANZEL(rufinamide) BRIVIACT (brivaracetam)	*Topiramate ER will be authorized after a thirty (30) day trial of topiramate IR.



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
divalproex divalproex ER divalproex sprinkle EPITOL (carbamazepine) GABITRIL (tiagabine) lamotrigine levetiracetam IR levetiracetam ER oxcarbazepine suspension and tablets topiramate IR topiramate ER* valproic acid VIMPAT(lacosamide) <sup>AP</sup> zonisamide	CARBATROL (carbamazepine) DEPAKENE (valproic acid) DEPAKOTE (divalproex) DEPAKOTE ER (divalproex) DEPAKOTE SPRINKLE (divalproex) EQUETRO (carbamazepine) FANATREX SUSPENSION (gabapentin) felbamate FELBATOL (felbamate) FYCOMPA (perampanel) KEPPRA (levetiracetam) KEPPRA XR (levetiracetam) LAMICTAL (lamotrigine) LAMICTAL ODT (lamotrigine) LAMICTAL XR (lamotrigine) LAMICTAL XR (lamotrigine) lamotrigine dose pack lamotrigine ER OXTELLAR XR (oxcarbazepine) POTIGA (ezogabine) QUDEXY XR (topiramate ER)** SABRIL (vigabatrin) SPRITAM (levetiracetam) STAVZOR (valproic acid) TEGRETOL (carbamazepine) TEGRETOL XR (carbamazepine) tiagabine TOPAMAX (topiramate) TRILEPTAL SUSPENSION and TABLETS (oxcarbazepine) TROKENDI XR (topiramate)** ZONEGRAN (zonisamide)	**Qudexy XR and Trokendi XR are only approvable on appeal.
	BARBITURATESAP	
phenobarbital primidone	MYSOLINE (primidone)	
	BENZODIAZEPINESAP	
clonazepam DIASTAT (diazepam rectal) diazepam tablets	clonazepam ODT diazepam rectal gel KLONOPIN (clonazepam) ONFI (clobazam)* ONFI SUSPENSION (clobazam)* VALIUM TABLETS (diazepam)	*Onfi shall be authorized as adjunctive therapy for treatment of Lennox-Gastaut Syndrome without further restrictions. Offlabel use requires an appeal to the Medical Director.



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLA	SS	
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
	HYDANTOINS <sup>AP</sup>		
DILANTIN (phenytoin sodium, extended) PEGANONE (ethotoin) phenytoin capsules, chewable tablets, suspension	DILANTIN INFATABS (phenytoin) PHENYTEK (phenytoin)		
	SUCCINIMIDES		
CELONTIN (methsuximide) ethosuximide capsules ethosuximide syrup	ZARONTIN (ethosuximide) capsules ZARONTIN (ethosuximide) syrup		
ANTIDEPRESSANTS, OTHER			
CLASS PA CRITERIA: See below for individu	ual sub-class criteria.		
	MAOIsAP		
	MARPLAN (isocarboxazid) NARDIL (phenelzine) PARNATE (tranylcypromine) phenelzine tranylcypromine	Patients stabilized on MAOI agents will be grandfathered.	
	SNRIS <sup>AP</sup>		
duloxetine capulses venlafaxine ER capsules	CYMBALTA (duloxetine) desvenlafaxine ER desvenlafaxine fumarate ER EFFEXOR XR (venlafaxine) FETZIMA (levomilnacipran) KHEDEZLA (desvenlafaxine) PRISTIQ (desvenlafaxine) venlafaxine IR VENLAFAXINE ER TABLETS (venlafaxine)	Non-preferred agents require separate thirty (30) day trials of a preferred agent in this sub-class <b>AND</b> an SSRI before they will be approved, unless one (1) of the exceptions on the PA form is present.	
	SECOND GENERATION NON-SSRI, OT	HERAP	
bupropion IR bupropion SR bupropion XL mirtazapine trazodone	APLENZIN (bupropion hbr) EMSAM (selegiline) FORFIVO XL (bupropion) nefazodone OLEPTRO ER (trazodone) REMERON (mirtazapine) TRINTELLIX (vortioxetine) VIIBRYD (vilazodone HCI) WELLBUTRIN (bupropion) WELLBUTRIN SR (bupropion) WELLBUTRIN XL (bupropion)	Non-preferred agents require separate thirty (30) day trials of a preferred agent in this sub-class <b>AND</b> an SSRI before they will be approved, unless one (1) of the exceptions on the PA form is present.	



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	S
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	SELECTED TCAs	
imipramine HCI	imipramine pamoate TOFRANIL (imipramine HCI) TOFRANIL PM (imipramine pamoate)	Non-preferred agents require a twelve (12) week trial of imipramine HCl before they will be approved, unless one (1) of the exceptions on the PA form is present.
ANTIDEPRESSANTS, SSRISAP		
CLASS PA CRITERIA: Non-preferred agents re exceptions on the PA form is present.	equire thirty (30) day trials of at least two (2) prefer	rred agents before they will be approved, unless one (1) of the
Upon hospital discharge, patients admitted with a continue that drug.	primary mental health diagnosis who have been stab	oilized on a non-preferred SSRI will receive an authorization to
citalopram escitalopram tablets fluoxetine capsules, solution fluvoxamine paroxetine sertraline	BRISDELLE (paroxetine) CELEXA (citalopram) escitalopram solution fluoxetine tablets fluvoxamine ER LEXAPRO (escitalopram) LUVOX CR (fluvoxamine) paroxetine 7.5 mg capsules paroxetine ER PAXIL (paroxetine) PAXIL CR (paroxetine) PEXEVA (paroxetine) PROZAC (fluoxetine) SARAFEM (fluoxetine) ZOLOFT (sertraline)	
ANTIEMETICSAP		
CLASS PA CRITERIA: See below for sub-class of	criteria.	
	5HT3 RECEPTOR BLOCKERS	
ondansetron ODT, solution, tablets	ANZEMET (dolasetron) granisetron GRANISOL (granisetron) ondansetron vials SANCUSO (granisetron) SUSTOL (granisetron) ZOFRAN (ondansetron) ZUPLENZ (ondansetron)	Non-preferred agents require a three (3) day trial of a preferred agent before they will be approved, unless one (1) of the exceptions on the PA form is present.
	CANNABINOIDS	10 (11)
	CESAMET (nabilone)* dronabinol**	*Cesamet will be authorized only for the treatment of nausea and vomiting associated with cancer chemotherapy for patients



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	MARINOL (dronabinol)** SYNDROS SOLUTION (dronabinol)	who have failed to respond adequately to three (3) day trials of conventional treatments such as promethazine or ondansetron and are eighteen (18) years of age or older.
		**Dronabinol will only be authorized for:  1. The treatment of anorexia associated with weight loss in patients with AIDS or cancer and unresponsive to megestrol or  2. The prophylaxis of chemotherapy induced nausea and vomiting unresponsive to three (3) day trials of ondansetron or promethazine for patients from eighteen (18) up to sixty-five (65) years of age.
	SUBSTANCE P ANTAGONISTS	
EMEND (aprepitant)	aprepitant VARUBI (rolapitant)	Non-preferred agents require a three (3) day trial of a preferred agent before they will be approved, unless one (1) of the exceptions on the PA form is present.
	COMBINATIONS	
	AKYNZEO (netupitant/palonosetron)	Non-preferred agents will only be approved on appeal.
ANTIFUNGALS, ORAL		
CLASS PA CRITERIA: Non-preferred agents wil	I only be authorized if one (1) of the exceptions on the	·
clotrimazole	ANCOBON (flucytosine)	*PA is required when limits are exceeded.
fluconazole* nystatin	CRESEMBA (isovuconazonium) <sup>CL**</sup> DIFLUCAN (fluconazole)	**Full PA criteria may be found on the PA Criteria page by
terbinafine <sup>CL</sup>	flucytosine	clicking the hyperlink.
	GRIFULVIN V TABLET (griseofulvin)	one.ung are rijperimuu
	griseofulvin***	***PA is not required for griseofulvin suspension for children up
	GRIS-PEG (griseofulvin)	to eighteen (18) years of age for the treatment of tinea capitis.
	itraconazole ketoconazole****	****Ketoconazole will be authorized if the following criteria are
	LAMISIL (terbinafine)	met:
	MYCELEX (clotrimazole)	Diagnosis of one of the following fungal infections:
	MYCOSTATIN Tablets (nystatin)	blastomycosis, coccidioidomycosis, histoplasmosis,
	NIZORAL (ketoconazole)	chromomycosis, or paracoccidioidomycosis and
	NOXAFIL (posaconazole)	2. Documented failure or intolerance of all other diagnosis-
	ONMEL (itraconazole) ORAVIG (miconazole)	appropriate antifungal therapies, i.e. itraconazole, fluconazole, flucytosine, etc <b>and</b>
	SPORANOX (itraconazole)	Baseline assessment of the liver status including alanine
	VFEND (voriconazole)	aminotransferase (ALT), aspartate aminotransferase
	voriconazole suspension	(AST), total bilirubin, alkaline phosphatase, prothrombin
	voriconazole tablets	time, and international normalized ratio (INR) before starting treatment <b>and</b>
		4. Weekly monitoring of serum ALT for the duration of



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	S
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		treatment (If ALT values increase to a level above the upper limit of normal or 30% above baseline, or if the patient develops symptoms of abnormal liver function, treatment should be interrupted and a full set of liver tests be obtained. Liver tests should be repeated to ensure normalization of values.) and  5. Assessment of all concomitant medications for potential adverse drug interactions with ketoconazole.  Ketoconazole will not be authorized for treatment for fungal infections of the skin and nails.
ANTIFUNGALS, TOPICALAP		
	mpoo is requested, a fourteen (14) day trial of one (1)	ats before they will be approved, unless one (1) of the exceptions preferred product (i.e. ketoconazole shampoo) is required.
	ANTIFUNGALS	
econazole ketoconazole cream, shampoo MENTAX (butenafine) miconazole (OTC) nystatin	CICLODAN (ciclopirox) ciclopirox ERTACZO (sertaconazole) EXELDERM (sulconazole) EXTINA (ketoconazole) JUBLIA (efinaconazole) ketoconazole foam KERYDIN (tavaborole) KETODAN (ketoconazole) LOPROX (ciclopirox) LUZU (luliconazole) MYCOSTATIN (nystatin) NAFTIN CREAM (naftifine) NAFTIN GEL (naftifine) NIZORAL (ketoconazole) OXISTAT (oxiconazole)* PEDIPIROX-4 (ciclopirox) PENLAC (ciclopirox) VUSION (miconazole/petrolatum/zinc oxide) XOLEGEL (ketoconazole)	*Oxistat cream will be authorized for children up to thirteen (13) years of age for tinea corporis, tinea cruris, tinea pedis, and tinea (pityriasis) versicolor.
clotrimazole/betamethasone	ANTIFUNGAL/STEROID COMBINATIO KETOCON PLUS (ketoconazole/hydrocortisone)	NO CALL
Sistemazio del Control de Control	LOTRISONE (clotrimazole/betamethasone) nystatin/triamcinolone	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS			
PREFERRED AGENTS PA CRITERIA			
ANTIHEMOPHILIA FACTOR AGENTS <sup>CL</sup> CLASS PA CRITERIA: All agents will require prior-authorization, and non-preferred agents require medical reasoning explaining why the need cannot be met using a preferred product.  All currently established regimens shall be grandfathered with documentation of adherence to therapy.			
	FACTOR VIII		
ALPHANATE HEMOFIL M HUMATE-P KOATE KOATE-DVI MONOCLATE-P NOVOEIGHT WILATE XYNTHA XYNTHA SOLOFUSE	ADVATE ADYNOVATE ELOCTATE KOGENATE FS KOVALTRY NUWIQ RECOMBINATE VONVENDI		
FACTOR IX			
ALPHANINE SD BEBULIN BENEFIX IXINITY MONONINE PROFILNINE RIXUBIS	ALPROLIX IDELVION		
<b>ANTIHYPERTENSIVES, SYMPATH</b>			
CLASS PA CRITERIA: Non-preferred agents r approved, unless one (1) of the exceptions on the CATAPRES-TTS (clonidine) clonidine tablets	equire thirty (30) day trials of each preferred unique choose PA form is present.  CATAPRES TABLETS (clonidine) clonidine patch NEXICLON XR (clonidine)	emical entity in the corresponding formulation before they will be	
ANTIHYPERURICEMICS			
	CLASS PA CRITERIA: Non-preferred agents require a thirty (30) day trial of one (1) of the preferred agents for the prevention of gouty arthritis attacks (colchicine/probenecid, probenecid, or allopurinol) before they will be approved, unless one (1) of the exceptions on the PA form is present.		
AUTIOARE ( . I. I ) *	ANTIMITOTICS		
MITIGARE (colchicine)*	colchicine capsules colchicine tablets COLCRYS (colchicine)	*In the case of acute gouty attacks, a ten (10) day supply (twenty (20) capsules) of Mitigare will be authorized per ninety (90) days.	



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	S		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA		
	ANTIMITOTIC-URICOSURIC COMBINATION			
colchicine/probenecid				
	URICOSURIC			
probenecid	ZURAMPIC (lesinurad)*	*Full PA criteria may be found on the PA Criteria page by clicking the hyperlink.		
	XANTHINE OXIDASE INHIBITORS			
allopurinol	ULORIC (febuxostat) ZYLOPRIM (allopurinol)			
	URICOSURIC – XANTHINE OXIDASE INHIE	BITORS		
	DUZALLO (allopurinol/lesinurad)	Non-preferred agents will only be approved on appeal.		
ANTIMIGRAINE AGENTS, OTHERAP				
CLASS PA CRITERIA: Non-preferred agents recapproved, unless one (1) of the exceptions on the		ty of the preferred Antimigraine Triptan Agents before they will be		
	CAMBIA (diclofenac)			
ANTIMIGRAINE AGENTS, TRIPTAN	IS <sup>AP</sup>			
<b>CLASS PA CRITERIA:</b> Non-preferred agents re exceptions on the PA form is present.	quire three (3) day trials of each preferred unique ch	nemical entity before they will be approved, unless one (1) of the		
	TRIPTANS			
naratriptan rizatriptan ODT rizatriptan tablet sumatriptan injection <sup>CL</sup> sumatriptan nasal spray sumatriptan tablets	almotriptan AMERGE (naratriptan) AXERT (almotriptan) eletriptan FROVA (frovatriptan) frovatriptan IMITREX INJECTION (sumatriptan) IMITREX NASAL SPRAY (sumatriptan) IMITREX tablets (sumatriptan) MAXALT MLT (rizatriptan) MAXALT (rizatriptan) ONZETRA XSAIL (sumatriptan)* RELPAX (eletriptan) SUMAVEL (sumatriptan) ZECUITY PATCH (sumatriptan) ZEMBRACE SYMTOUCH (sumatriptan) zolmitriptan zolmitriptan ODT	*In addition to the Class Criteria: Onzetra Xsail requires three (3) day trials of each preferred oral, nasal and injectable forms of sumatriptan.		



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	SS
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	ZOMIG (zolmitriptan) ZOMIG ZMT (zolmitriptan) TRIPTAN COMBINATIONS	
	TREXIMET (sumatriptan/naproxen sodium)	
ANTIPARASITICS, TOPICAL <sup>AP</sup>		
CLASS PA CRITERIA: Non-preferred agents rec (1) of the exceptions on the PA form is present.	uire trials of each preferred agent (which are age and	d weight appropriate) before they will be approved, unless one
NATROBA (spinosad) permethrin 5% cream permethrin 1% lotion (OTC) pyrethrins-piperonyl butoxide OTC SKLICE (ivermectin)	EURAX (crotamiton) LICE EGG REMOVER OTC (benzalkonium chloride) lindane malathion OVIDE (malathion) spinosad	
ANTIPARKINSON'S AGENTS		
<b>CLASS PA CRITERIA:</b> Patients starting therapy a non-preferred agent will be authorized.	_	gy to all preferred agents in the corresponding sub-class, before
	ANTICHOLINERGICS	
benztropine trihexyphenidyl	COGENTIN (benztropine)	
	COMT INHIBITORS	
	COMTAN (entacapone) entacapone TASMAR (tolcapone)	COMT Inhibitor agents will only be approved as add-on therapy to a levodopa-containing regimen for treatment of documented motor complications.
	DOPAMINE AGONISTS	
pramipexole ropinirole	MIRAPEX (pramipexole) MIRAPEX ER (pramipexole)* NEUPRO (rotigotine) pramipexole ER REQUIP (ropinirole) REQUIP XL (ropinirole)* ropinirole ER	*Mirapex ER and Requip XL will be authorized for a diagnosis of Parkinsonism without a trial of preferred agents.
	OTHER ANTIPARKINSON'S AGENT	
amantadine*AP bromocriptine carbidopa/levodopa levodopa/carbidopa/entacapone selegiline	AZILECT (rasagiline) carbidopa ELDEPRYL (selegiline) levodopa/carbidopa ODT LODOSYN (carbidopa) PARCOPA (levodopa/carbidopa) PARLODEL (bromocriptine)	*Amantadine will not be authorized for the treatment or prophylaxis of influenza.



**EFFECTIVE** 07/01/2018 Version 2018.3e

This is not an all-inclusive list of available covered drugs and includes only managed categories. Refer to cover page for complete list of rules governing this PDL.

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	rasagiline RYTARY (levodopa/carbidopa) SINEMET (levodopa/carbidopa) STALEVO (levodopa/carbidopa/entacapone) XADAGO (safinamide) ZELAPAR (selegiline)	
ANTIPSORIATICS, TOPICAL		
<b>CLASS PA CRITERIA:</b> Non-preferred agents require thirty (30) day trials of two (2) preferred unique chemical entities before they will be approved, unless one (1) of the exceptions on the PA form is present.		

TACLONEX OINT (calcipotriene/	calcipotriene cream	
betamethasone)	calcipotriene ointment	
TAZORAC (tazarotene)	calcipotriene solution	
VECTICAL (calcitriol)	calcipotriene/betamethasone ointment	
	CALCITRENE (calcipotriene)	
	calcitriol	
	DOVONEX (calcipotriene)	
	ENSTILAR (calcipotriene/betamethasone)	
	SORILUX (calcipotriene)	
	tazarotene cream (tazarotene)	
	, ,	

### **ANTIPSYCHOTICS, ATYPICAL**

CLASS PA CRITERIA: All antipsychotic agents require prior authorization for children up to eighteen (18) years of age. All PA requests for antipsychotics for children 6 years of age and younger will be reviewed by Medicaid's consultant psychiatrist.

Non-preferred agents require fourteen (14) day trials of three (3) preferred agents, including the generic formulation of the requested agent (if available), before they will be approved unless one (1) of the exceptions on the PA form is present.

Upon discharge, a hospitalized patient stabilized on a non-preferred agent may receive authorization to continue this drug for labeled indications and at FDA recommended dosages. For off-label indications or dosages, a thirty (30) day prior-authorization shall be granted pending BMS review.

	SINGLE INGREDIENT	
ABILIFY MAINTENA (aripiprazole) <sup>CL</sup>	ABILIFY TABLETS (aripiprazole)	In addition to class criteria:
aripiprazole tablets	ADASUVE (loxapine)	
ARISTADA (aripiprazole) <sup>CL</sup>	clozapine ODT	*Invega Trinza will be authorized after four months' treatment
clozapine	CLOZARIL (clozapine)	with Invega Sustenna
INVEGA SUSTENNA (paliperidone)CL	FANAPT (iloperidone)	_
INVEGA TRINZA (paliperidone)* CL	FAZACLO (clozapine)	**Quetiapine 25 mg will be authorized:
olanzapine	GEODON (ziprasidone)	<ol> <li>For a diagnosis of schizophrenia or</li> </ol>
olanzapine ODT	GEODON IM (ziprasidone)	<ol><li>For a diagnosis of bipolar disorder or</li></ol>
quetiapine** AP for the 25 mg Tablet Only	INVEGA ER (paliperidone)	3. When prescribed concurrently with other strengths of
quetiapine ER	LATUDA (lurasidone)*** AP	Seroquel in order to achieve therapeutic treatment



EFFECTIVE 07/01/2018 Version 2018.3e

This is not an all-inclusive list of available covered drugs and includes only managed categories. Refer to cover page for complete list of rules governing this PDL.

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
RISPERDAL CONSTA (risperidone) <sup>CL</sup> risperidone ziprasidone	NUPLAZID (pimavanserin) **** olanzapine IMCL paliperidone ER REXULTI (brexipiprazole) RISPERDAL (risperidone) SAPHRIS (asenapine) SEROQUEL (quetiapine) SEROQUEL XR (quetiapine) VERSACLOZ (clozapine) VRAYLAR (capriprazine) VRAYLAR DOSE PAK (capriprazine) ZYPREXA (olanzapine) ZYPREXA IM (olanzapine) CL ZYPREXA RELPREVV (olanzapine) ATYPICAL ANTIPSYCHOTIC/SSRI COMBIN olanzapine/fluoxetine SYMBYAX (olanzapine/fluoxetine)	levels.  Quetiapine 25 mg will not be authorized for use as a sedative hypnotic.  ***For the indication of bipolar depression only, prior authorization of Latuda requires a 14-day trial of either quetiapine OR a combination of olanzapine + fluoxetine. All other indications follow class criteria. Patients already stabilized on Latuda shall be grandfathered.  ****Nuplazid will only be authorized for the treatment of Parkinson Disease Induced Psychosis after documented treatment failure with quetiapine.  ATIONS

#### **ANTIRETROVIRALS**

**CLASS PA CRITERIA:** Non-preferred drugs require medical reasoning beyond convenience or enhanced compliance as to why the clinical need cannot be met with a preferred agent or combination of preferred agents. <u>NOTE</u>: Regimens consisting of preferred agents will result in no more than one additional unit per day over equivalent regimens composed of non-preferred agents. Patients already on a non-preferred regimen shall be grandfathered.

#### INTEGRASE STRAND TRANSFER INHIBITORS ISENTRESS (raltegravir potassium) TIVICAY (dolutegravir sodium) VITEKTA (elvitegravir) **NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (NRTI)** abacavir sulfate tablet abacavir sulfate solution<sup>NR</sup> didanosine DR capsule EPIVIR TABLET (lamivudine) EMTRIVA (emtricitabine) RETROVIR (zidovudine) **EPIVIR SOLUTION (lamivudine)** VIDEX EC (didanosine) lamivudine ZERIT (stavudine) stavudine ZIAGEN TABLET (abacavir sulfate) VIDEX SOLUTION (didanosine) VIREAD (tenofovir disoproxil fumarate) ZIAGEN SOLUTION (abacavir sulfate) zidovudine NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITOR (NNRTI) EDURANT (rilpivirine) INTELENCE (etravirine) SUSTIVA (efavirenz) nevirapine nevirapine ER



EFFECTIVE 07/01/2018 Version 2018.3e

TYBOST (cobicistat)	NON-PREFERRED AGENTS RESCRIPTOR (delavirdine mesylate) (IRAMUNE ER 24H (nevirapine) (IRAMUNE SUSPENSION (nevirapine) PHARMACOENHANCER – CYTOCHROME P450 PROTEASE INHIBITORS (PEPTIDIC) CRIXIVAN (indinavir)	
R V V V F TYBOST (cobicistat)	RESCRIPTOR (delavirdine mesylate) (IRAMUNE ER 24H (nevirapine) (IRAMUNE SUSPENSION (nevirapine) PHARMACOENHANCER – CYTOCHROME P450 PROTEASE INHIBITORS (PEPTIDIC) CRIXIVAN (indinavir)	) INHIBITOR
TYBOST (cobicistat)	(IRAMUNE ER 24H (nevirapine) (IRAMUNE SUSPENSION (nevirapine) PHARMACOENHANCER – CYTOCHROME P450 PROTEASE INHIBITORS (PEPTIDIC) CRIXIVAN (indinavir)	
TYBOST (cobicistat)	PHARMACOENHANCER – CYTOCHROME P450  PROTEASE INHIBITORS (PEPTIDIC)  CRIXIVAN (indinavir)	
TYBOST (cobicistat)	PHARMACOENHANCER – CYTOCHROME P450  PROTEASE INHIBITORS (PEPTIDIC)  CRIXIVAN (indinavir)	
, ,	CRIXIVAN (indinavir)	
EVOTAZ (atazanavir/cobicistat) C	CRIXIVAN (indinavir)	
EVOTAZ (atazanavir/cobicistat)	CRIXIVAN (indinavir)	
NORVIR (ritonavir)	NVIRASE (saquinavir mesylate)	
	<mark>osamprenavir</mark>	
	EXIVA (fosamprenavir)	
V	IRACEPT (nelfinavir mesylate)	10\
PREZISTA (darunavir ethanolate) A	PROTEASE INHIBITORS (NON-PEPTID	nc)
	PTIVUS (tipranavir) REZCOBIX (darunavir/cobicistat)	
	NTRY INHIBITORS - CCR5 CO-RECEPTOR AN	TACOMICTO
	ELZENTRY (maraviroc)	TAGONISTS
3	ENTRY (MAIAVIOC)  ENTRY INHIBITORS – FUSION INHIBITO	nne
_		JRS
-	UZEON (enfuvirtide)  COMBINATION PRODUCTS - NRTIS	
EPZICOM (abacavir/lamivudine) al	bacavir/lamivudine <sup>NR</sup>	
	bacavir/lamivudine/zidovudine	
	COMBIVIR (lamivudine/zidovudine)	
	RIZIVIR (abacavir/lamivudine/zidovudine)	
	TTZTVTT (abacavii/laitiivuulite/2luovuulite)	
COMBINATION PRODUCT	TS – INTEGRASE STRAND TRANSFER INHIBIT	ORS & NUCLEOSIDE ANALOG RTIS
BIKTARVY (bictegravir/emtricitabine/tenofovir		
alafenamide)	ATION PRODUCTS - NUCLEOSIDE & NUCLEO	TIDE ANALOG RTIS
DESCOVY (emtricitabine/tenofovir)	ATION I NODUCTO - NOCELOGIDE & NOCELO	TIDE ANALOG KTIS
TRUVADA (emtricitabine/tenofovir)		
	DUCTS - NUCLEOSIDE & NUCLEOTIDE ANALO	OGS & INTEGRASE INHIBITORS
GENVOYA S	TRIBILD	*Stribild requires medical reasoning beyond convenience or
(elvitegravir/cobicistat/emtricitabine/tenofovir)	(elvitegravir/cobicistat/emtricitabine/tenofovir)*	enhanced compliance as to why the medical need cannot be
T	RIUMEQ (abacavir/lamivudine/ dolutegravir)**	met with the the preferred agent Genvoya.
		** <del>*</del> · · · · · · · · · · · ·
		**Triumeq requires medical reasoning beyond convenience or enhanced compliance as to why the medical need cannot be
		met with the preferred agents Epzicom and Tivicay.



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
COMBINATION P	RODUCTS - NUCLEOSIDE & NUCLEOTIDE ANAL	OGS & NON-NUCLEOSIDE RTIs
ATRIPLA (efavirenz/emtricitabine/tenofovir) ODEFSEY (emtricitabine/rilpivirine/tenofovir)	COMPLERA (emtricitabine/rilpivirine/tenofovir)*	*Complera requires medical reasoning beyond convenience or enhanced compliance as to why the medical need cannot be met with the preferred agents Truvada and Edurant.
	COMBINATION PRODUCTS - PROTEASE IN	HIBITORS
KALETRA (lopinavir/ritonavir)	lopinavir/ritonavir	
ANTIVIRALS, ORAL		
<b>CLASS PA CRITERIA:</b> Non-preferred agents rethe exceptions on the PA form is present.	quire five (5) day trials of each preferred agent in the	same sub-class before they will be approved, unless one (1) of
	ANTI HERPES	
acyclovir	famciclovir	
valacyclovir	FAMVIR (famciclovir) SITAVIG (acyclovir) VALTREX ZOVIRAX (acyclovir)	
	ANTI-INFLUENZA	
RELENZA (zanamivir) TAMIFLU (oseltamivir)	FLUMADINE (rimantadine) oseltamivir rimantadine	In addition to the Class Criteria: The anti-influenza agents will be authorized only for a diagnosis of influenza.
ANTIVIRALS, TOPICALAP		
·	quire a five (5) day trial of the preferred agent before t	they will be approved, unless one (1) of the exceptions on the PA
ZOVIRAX CREAM (acyclovir)	ABREVA (docosanol) acyclovir ointment DENAVIR (penciclovir) ZOVIRAX OINTMENT (acyclovir)	
BETA BLOCKERSAP		
	quire fourteen (14) day trials of three (3) chemically die approved, unless one (1) of the exceptions on the Pa	istinct preferred agents, including the generic formulation of the A form is present.
BETA BLOCKERS		
acebutolol atenolol betaxolol bisoprolol	BETAPACE (sotalol) BYSTOLIC (nebivolol) HEMANGEOL (propranolol)* INDERAL LA (propranolol)	*Hemangeol will be authorized for the treatment of proliferating infantile hemangioma requiring systemic therapy.  **Propranolol ER shall be authorized for patients with a
CORGARD (nadolol) metoprolol	INDERAL XL (propranolol) INNOPRAN XL (propranolol)	diagnosis of migraines. Existing users will be grandfathered for use in migraine prophylaxis.



EFFECTIVE 07/01/2018 Version 2018.3e

This is not an all-inclusive list of available covered drugs and includes only managed categories. Refer to cover page for complete list of rules governing this PDL.

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
metoprolol ER pindolol propranolol sotalol timolol	KERLONE (betaxolol) LEVATOL (penbutolol) LOPRESSOR (metoprolol) nadolol propranolol ER** SECTRAL (acebutolol) TENORMIN (atenolol) TOPROL XL (metoprolol) ZEBETA (bisoprolol)	
	BETA BLOCKER/DIURETIC COMBINATION	DRUGS
atenolol/chlorthalidone bisoprolol/HCTZ metoprolol/HCTZ propranolol/HCTZ	CORZIDE (nadolol/bendroflumethiazide) DUTOPROL (metoprolol ER/HCTZ ER) LOPRESSOR HCT (metoprolol/HCTZ) metoprolol/HCTZ ER nadolol/bendroflumethiazide TENORETIC (atenolol/chlorthalidone) ZIAC (bisoprolol/HCTZ)	
BETA- AND ALPHA-BLOCKERS		
carvedilol labetalol	COREG (carvedilol) COREG CR (carvedilol) TRANDATE (labetalol)	
BLADDER RELAXANT PREPARATIONS <sup>AP</sup>		
CLASS DA CRITERIA. Non preferred agents require thirty (20) day triple of each chemically distinct preferred agent before they will be approved upless one (4) of the		

CLASS PA CRITERIA: Non-preferred agents require thirty (30) day trials of each chemically distinct preferred agent before they will be approved, unless one (1) of the exceptions on the PA form is present

oxybutynin IR oxybutynin ER TOVIAZ (fesoterodine)	DETROL (tolterodine) DETROL LA (tolterodine) DITROPAN XL (oxybutynin) ENABLEX (darifenacin) flavoxate GELNIQUE (oxybutynin) MYRBETRIQ (mirabegron) OXYTROL (oxybutynin) SANCTURA (trospium) SANCTURA XR (trospium) tolterodine tolterodine ER trospium trospium ER VESICARE (solifenacin)	
---------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
BONE RESORPTION SUPPRES	SION AND RELATED AGENTS		
CLASS PA CRITERIA: See below for class	criteria.		
	BISPHOSPHONATES		
alendronate tablets ibandronate	ACTONEL (risedronate) ACTONEL WITH CALCIUM (risedronate/ calcium) alendronate solution ATELVIA (risedronate) BINOSTO (alendronate) BONIVA (ibandronate) DIDRONEL (etidronate) etidronate FOSAMAX TABLETS (alendronate) FOSAMAX PLUS D (alendronate/vitamin D) risedronate	Non-preferred agents require thirty (30) day trials of <b>each</b> preferred Bisphosphonate agent before they will be approved, unless one (1) of the exceptions on the PA form is present.	
	OTHER BONE RESORPTION SUPPRESSION AND		
	calcitonin EVISTA (raloxifene)* FORTEO (teriparatide) FORTICAL (calcitonin) MIACALCIN (calcitonin) raloxifene* TYMLOS (abaloparatide)	Non-preferred agents require a thirty (30) day trial of a preferred Bisphosphonate agent before they will be approved, unless one (1) of the exceptions on the PA form is present.  *Raloxifene generic will be authorized for postmenopausal women with osteoporosis or at high risk for invasive breast cancer.	
BPH TREATMENTS			
	s require thirty (30) day trials of at least two (2) chemically will be approved, unless one (1) of the exceptions on	ally distinct preferred agents, including the generic formulation of the PA form is present.	
	5-ALPHA-REDUCTASE (5AR) INHIBITORS AN	D PDE-5 AGENTS	
finasteride	AVODART (dutasteride) CIALIS 5 mg (tadalafil) dutasteride PROSCAR (finasteride)		
alfuzosin	ALPHA BLOCKERS  CARDURA (doxazosin)		
artuzosin doxazosin tamsulosin terazosin	CARDURA (doxazosin) CARDURA XL (doxazosin) FLOMAX (tamsulosin) HYTRIN (terazosin) RAPAFLO (silodosin) UROXATRAL (alfuzosin)		



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
5-ALI	PHA-REDUCTASE (5AR) INHIBITORS/ALPHA BLO	
	dutasteride/tamsulosin JALYN (dutasteride/tamsulosin)	<b>Substitute for Class Criteria</b> : Concurrent thirty (30) day trials of dutasteride and tamsulosin are required before the non-preferred agent will be authorized.
BRONCHODILATORS, BETA AGON	IIST <sup>ap</sup>	
<b>CLASS PA CRITERIA:</b> Non-preferred agents red the exceptions on the PA form is present.	uire thirty (30) day trials of each chemically distinct p	preferred agent in their corresponding sub-class unless one (1) of
	INHALATION SOLUTION	
albuterol	BROVANA (arformoterol) levalbuterol metaproterenol PERFOROMIST (formoterol) XOPENEX (levalbuterol)*	*Xopenex Inhalation Solution will be authorized for twelve (12) months for a diagnosis of asthma or COPD for patients on concurrent asthma controller therapy (either oral or inhaled) with documentation of failure on a trial of albuterol or documented intolerance of albuterol, or for concurrent diagnosis of heart disease.
	INHALERS, LONG-ACTING	
FORADIL (formoterol) SEREVENT (salmeterol)	ARCAPTA (indacaterol maleate) STRIVERDI RESPIMAT (olodaterol)	
DDOAID LIEA (-lb.:/l)	INHALERS, SHORT-ACTING	
PROAIR HFA (albuterol) PROVENTIL HFA (albuterol)	MAXAIR (pirbuterol) PROAIR RESPICLICK (albuterol) VENTOLIN HFA (albuterol) XOPENEX HFA (levalbuterol)	
	ORAL	
albuterol ER albuterol IR terbutaline	metaproterenol VOSPIRE ER (albuterol)	
CALCIUM CHANNEL BLOCKERSAP		
CLASS PA CRITERIA: Non-preferred agents recunless one (1) of the exceptions on the PA form is		within the corresponding sub-class before they will be approved,
	LONG-ACTING	
amlodipine diltiazem ER felodipine ER nifedipine ER verapamil ER	ADALAT CC (nifedipine) CALAN SR (verapamil) CARDENE SR (nicardipine) CARDIZEM CD, LA (diltiazem) COVERA-HS (verapamil) diltiazem LA DYNACIRC CR (isradipine) ISOPTIN SR (verapamil) MATZIM LA (diltiazem) nisoldipine	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS			
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
	NORVASC (amlodipine) PLENDIL (felodipine) PROCARDIA XL (nifedipine) SULAR (nisoldipine) TIAZAC (diltiazem) verapamil ER PM VERELAN/VERELAN PM (verapamil)		
	SHORT-ACTING		
diltiazem verapamil	CALAN (verapamil) CARDIZEM (diltiazem) isradipine nicardipine nifedipine nimodipine NIMOTOP (nimodipine) NYMALIZE SOLUTION (nimodipine) PROCARDIA (nifedipine)		
CEPHALOSPORINS AND RELATED			
one (1) of the exceptions on the PA form is preser	CLASS PA CRITERIA: Non-preferred agents require a five (5) day trial of a preferred agent within the corresponding sub-class before they will be approved, unless one (1) of the exceptions on the PA form is present.		
	TAMS AND BETA LACTAM/BETA-LACTAMASE IN	HIBITOR COMBINATIONS	
amoxicillin/clavulanate IR	amoxicillin/clavulanate ER AUGMENTIN (amoxicillin/clavulanate) AUGMENTIN XR (amoxicillin/clavulanate) MOXATAG (amoxicillin)		
	CEPHALOSPORINS		
cefaclor capsule cefadroxil capsule, tablet cefdinir cefuroxime tablet cephalexin capsule, suspension	CEDAX (ceftibuten) cefaclor suspension cefaclor ER tablet cefadroxil suspension cefditoren cefpodoxime cefprozil ceftibuten capsule, suspension CEFTIN (cefuroxime) cefuroxime suspension cephalexin tablet KEFLEX (cephalexin) OMNICEF (cefdinir) RANICLOR (cefaclor) SPECTRACEF (cefditoren) SUPRAX (cefixime)		



EFFECTIVE 07/01/2018 Version 2018.3e

CLASS PA CRITERIA: Non-preferred agents require a sixty (60) day trial of one preferred agent from the corresponding sub-class before they will be approved, unless one (1) of the exceptions on the PA form is present.  ANTICHOLINERGICAP    Pratropium nebulizer solution   SPIRIVA (tiotropium)   SEEBRI NEOHALER(glycoppyrolate)   SPIRIVA (tiotropium)   SEEBRI NEOHALER(glycoppyrolate)   SPIRIVA (RESPIMAT (diotropium)	THERAPEUTIC DRUG CLASS		
CLASS PA CRITERIA: Non-preferred agents require a sixty (60) day trial of one preferred agent from the corresponding sub-class before they will be approved, unless one (1) of the exceptions on the PA form is present.  ANTICHOLINERGICAP  ATROVENT HFA ((prestropium)  ARROVENT HFA ((prestropium)  SPIRIVA (tiotropium)  ARROVENT HFA ((prestropium)  SPIRIVA (tiotropium)  ANTICHOLINERGIC-BETA AGONIST COMBINATIONSAP  ANTICHOLINERGIC-BETA AGONIST COMBINATIONSAP  ANTICHOLINERGIC-BETA AGONIST COMBINATIONSAP  ANTICHOLINERGIC-BETA AGONIST COMBINATIONSAP  ANTICHOLINERGIC-BETA AGONIST-GLUCOCORTICOID COMBINATIONS  Titelogy (albuterol/ipratropium)  STIDLTO RESPIMAT ((albuterol/ipratropium))  STIDLE SPILIPSI ((ulicasone/umedidinium/vilanterol))  Titelogy (Ellipsi (ulicasone/umedidinium/vilanterol))  Titelogy (Ellipsi (ulicasone/umedidinium/vilanterol))  **STIDLTO RESPIMAT ((albuterol/ipratropium))  STIDLE SPILIPSI (ulicasone/umedidinium/vilanterol))  Titelogy (Ellipsi (ulicasone/umedidinium/vilanterol))  **STIDLTO RESPIMAT ((albuterol/ipratropium))  **DALIRESP (roflumillast)**  **Daliresp will be authorized if the following criteria are met:  1. Patient is forty (40) years of age or older and  2. Diagnosis of severe chronic obstructive pulmonary disease (COPD) associated with chronic bronchitis and multiple exacerbations requiring systemic glucocorticodis in the preceding six (6) months and  3. Concurrent therapy with an inhaled corticosteroid and  4. No evidence of moderate to severe liver impairment (Child-Pugh Class B or C) and  5. No concurrent use with strong cytochrome P450 inducers (rifampicin, phenobarbital, carbamazepine or phenytoin)  CYTOKINE & CAM ANTAGONISTSc.*  CLASS PA CRITERIA: Non-preferred agents require ninety (90) day trial of Cosentyx will also be required.  ANTI-TNFS	PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ATROVENT HFA (ipratropium) SPIRIVA (tiotropium) SPIRIVA (tiotropium) SPIRIVA (tiotropium) SPIRIVA (tiotropium) SPIRIVA (tiotropium) SPIRIVA (tiotropium) NCRUSE ELLIPTA (umeclidinium) SEEBI NEOHALER (glycopyrolate) SPIRIVA RESPIMAT (diotropium) TUDORAZ (actidinium) ANTICHOLINERGIC-BETA AGONIST COMBINATIONS  ANTICHOLINERGIC-BETA AGONIST COMBINATIONS  ANTICHOLINERGIC-BETA AGONIST COMBINATIONS  TOUR ESPIMAT (idiotropium) STIOL TO RESPIMAT (idiotropium) STIOL TO RESPIMAT (idiotropium) STIOL (idiotropium) S	COPD AGENTS		
ATROVENT HFA (ipratropium)   ATROVENT HFA (ipratropium)   INCRUSE ELLIPTA (umeclidinium)   SEBRI NEOHALER(glycopyrrolate)   SPIRIVA RESPIMAT (idotropium)   TUDORZA (aclidinium)   TUDORZA (aclidinium)   ANTICHOLINERGIC-BETA AGONIST COMBINATIONSA*			from the corresponding sub-class before they will be approved,
SPIRIVA (tiotropium)  INCRUSE ELLIPTĂ (umeclidinium) SEER NECHALER (glycopyrrolate) SPIRIVA RESPIMAT (tiotropium) TUDORZA (acidinium) ANTCHOLINERGIC-BETA AGONIST COMBINATIONS^*  ANORO ELLIPTĂ (umeclidinium/vilanterol) COMBIVENT RESPIMAT (tiotropium) DUONEB (albuterol/ipratropium) STIOLTO RESPIMAT (tiotropium) OUNEB (albuterol/ipratropium) STIOLTO RESPIMAT (tiotropium) STIOLTO RESPIMAT (tiotrop			
ANORO ELLIPTA (umeclidinium/vilanterol) COMBIVENT RESPIMAT (albuterol/ipratropium) DUONEB (albuterol/ipratropium) DUONEB (albuterol/ipratropium) STIOLTO RESPIMAT (flotropium/olodaterol)* UTIBRON (indacaterol/glycopyrrolate)  ANTICHOLINERGIC-BETA AGONIST-GLUCOCORTICOID COMBINATIONS  Trelegy Ellipta (fluticasone/umeclidinium/vilanterol)*  PDE4 INHIBITOR  DALIRESP (roflumilast)*  DALIRESP (roflumilast)*  *Daliresp will be authorized if the following criteria are met: 1. Patient is forty (40) years of age or older and 2. Diagnosis of severe chronic obstructive pulmonary disease (COPD) associated with chronic bronchitis and multiple exacerbations requiring systemic glucocorticoids in the preceding six (6) months and 3. Concurrent therapy with an inhaled corticosteroid and long-acting bronchodilator and evidence of compliance and 4. No evidence of moderate to severe liver impairment (Child-Pugh Class B or C) and 5. No concurrent use with strong cytochrome P450 inducers (rifampicin, phenobarbital, carbamazepine or phenytoin)  CYTOKINE & CAM ANTAGONISTS <sup>CL</sup> CLASS PA CRITERIA: Non-preferred agents require ninety (90) day trial of Cosentyx will also be required.  ANTI-TNFS	Ipratropium nebulizer solution SPIRIVA (tiotropium)	INCRUSE ELLIPTA (umeclidinium) SEEBRI NEOHALER(glycopyrrolate) SPIRIVA RESPIMAT (tiotropium)	
BEVESPI (glycopyrrolate/formoterol)  COMBIVENT RESPIMAT (albuterol/ipratropium) DUONEB (albuterol/ipratropium) STIOLTO RESPIMAT (iotropium/olodaterol)* UTIBRON (indacaterol/glycopyrrolate)  ANTICHOLINERGIC-BETA AGONIST-GLUCOCORTICOID COMBINATIONS  Trelegy Ellipta (fluticasone/umeclidinium/vilanterol)*  PDE4 INHIBITOR  DALIRESP (roflumilast)*  *Daliresp will be authorized if the following criteria are met: 1. Patient is forty (40) years of age or older and 2. Diagnosis of severe chronic obstructive pulmonary disease (COPD) associated with chronic bronchitis and multiple exacerbations requiring systemic glucocorticoids in the preceding six (6) months and 3. Concurrent therapy with an inhaled corticosteroid and long-acting bronchodilator and evidence of compliance and 4. No evidence of moderate to severe liver impairment (Child-Pugh Class B or C) and 5. No concurrent use with strong cytochrome P450 inducers (rifampicin, phenobarbital, carbamazepine or phenytoin)  CYTOKINE & CAM ANTAGONISTS <sup>CL</sup> CLASS PA CRITERIA: Non-preferred agents require ninety (90) day trials of both Humira and Enbrel unless one (1) of the exceptions on the PA form is present. For FDA-approved indications, an additional ninety (90) day trials of Cosentyx will also be required.  ANTI-TNFS		ANTICHOLINERGIC-BETA AGONIST COMBIN	NATIONSAP
Trelegy Ellipta (fluticasone/umeclidinium/vilanterol)*  PDE4 INHIBITOR  DALIRESP (roflumilast)*  *Daliresp will be authorized if the following criteria are met:  1. Patient is forty (40) years of age or older and  2. Diagnosis of severe chronic obstructive pulmonary disease (COPD) associated with chronic bronchitis and multiple exacerbations requiring systemic glucocorticoids in the preceding six (6) months and  3. Concurrent therapy with an inhaled corticosteroid and long-acting bronchodilator and evidence of compliance and  4. No evidence of moderate to severe liver impairment (Child-Pugh Class B or C) and  5. No concurrent use with strong cytochrome P450 inducers (rifampicin, phenobarbital, carbamazepine or phenytoin)  CYTOKINE & CAM ANTAGONISTS <sup>CL</sup> CLASS PA CRITERIA: Non-preferred agents require ninety (90) day trials of both Humira and Enbrel unless one (1) of the exceptions on the PA form is present. For FDA-approved indications, an additional ninety (90) day trial of Cosentyx will also be required.  ANTI-TNFS	albuterol/ipratropium nebulizer solution BEVESPI (glycopyrrolate/formoterol)	COMBIVENT RESPIMAT (albuterol/ipratropium) DUONEB (albuterol/ipratropium) STIOLTO RESPIMAT (tiotropium/olodaterol)* UTIBRON (indacaterol/glycopyrrolate)	sixty (60) day trial of Anoro Ellipta.
PDE4 INHIBITOR  DALIRESP (roflumilast)*  *Daliresp will be authorized if the following criteria are met:  1. Patient is forty (40) years of age or older and  2. Diagnosis of severe chronic obstructive pulmonary disease (COPD) associated with chronic bronchitis and multiple exacerbations requiring systemic glucocorticoids in the preceding six (6) months and  3. Concurrent therapy with an inhaled corticosteroid and long-acting bronchodilator and evidence of compliance and  4. No evidence of moderate to severe liver impairment (Child-Pugh Class B or C) and  5. No concurrent use with strong cytochrome P450 inducers (rifampicin, phenobarbital, carbamazepine or phenytoin)  CYTOKINE & CAM ANTAGONISTS <sup>CL</sup> CLASS PA CRITERIA: Non-preferred agents require ninety (90) day trials of both Humira and Enbrel unless one (1) of the exceptions on the PA form is present. For FDA-approved indications, an additional ninety (90) day trial of Cosentyx will also be required.  ANTI-TNFS	ANT ANT	ICHOLINERGIC-BETA AGONIST-GLUCOCORTICO	OID COMBINATIONS
DALIRESP (roflumilast)*  *Daliresp will be authorized if the following criteria are met:  1. Patient is forty (40) years of age or older and  2. Diagnosis of severe chronic obstructive pulmonary disease (COPD) associated with chronic bronchitis and multiple exacerbations requiring systemic glucocorticoids in the preceding six (6) months and  3. Concurrent therapy with an inhaled corticosteroid and long-acting bronchodilator and evidence of compliance and  4. No evidence of moderate to severe liver impairment (Child-Pugh Class B or C) and  5. No concurrent use with strong cytochrome P450 inducers (rifampicin, phenobarbital, carbamazepine or phenytoin)  CYTOKINE & CAM ANTAGONISTS <sup>CL</sup> CLASS PA CRITERIA: Non-preferred agents require ninety (90) day trials of both Humira and Enbrel unless one (1) of the exceptions on the PA form is present. For FDA-approved indications, an additional ninety (90) day trial of Cosentyx will also be required.  ANTI-TNFs		Trelegy Ellipta (fluticasone/umeclidinium/vilanterol)*	
1. Patient is forty (40) years of age or older and 2. Diagnosis of severe chronic obstructive pulmonary disease (COPD) associated with chronic bronchitis and multiple exacerbations requiring systemic glucocorticoids in the preceding six (6) months and 3. Concurrent therapy with an inhaled corticosteroid and long-acting bronchodilator and evidence of compliance and 4. No evidence of moderate to severe liver impairment (Child-Pugh Class B or C) and 5. No concurrent use with strong cytochrome P450 inducers (rifampicin, phenobarbital, carbamazepine or phenytoin)  CYTOKINE & CAM ANTAGONISTS <sup>CL</sup> CLASS PA CRITERIA: Non-preferred agents require ninety (90) day trials of both Humira and Enbrel unless one (1) of the exceptions on the PA form is present. For FDA-approved indications, an additional ninety (90) day trial of Cosentyx will also be required.  ANTI-TNFS		PDE4 INHIBITOR	
CLASS PA CRITERIA: Non-preferred agents require ninety (90) day trials of both Humira and Enbrel unless one (1) of the exceptions on the PA form is present. For FDA-approved indications, an additional ninety (90) day trial of Cosentyx will also be required.  ANTI-TNFs			<ol> <li>Patient is forty (40) years of age or older and</li> <li>Diagnosis of severe chronic obstructive pulmonary disease (COPD) associated with chronic bronchitis and multiple exacerbations requiring systemic glucocorticoids in the preceding six (6) months and</li> <li>Concurrent therapy with an inhaled corticosteroid and long-acting bronchodilator and evidence of compliance and</li> <li>No evidence of moderate to severe liver impairment (Child-Pugh Class B or C) and</li> <li>No concurrent use with strong cytochrome P450 inducers (rifampicin, phenobarbital, carbamazepine or</li> </ol>
FDA-approved indications, an additional ninety (90) day trial of Cosentyx will also be required.  ANTI-TNFs	CYTOKINE & CAM ANTAGONISTS	CL	
	<b>CLASS PA CRITERIA:</b> Non-preferred agents require ninety (90) day trials of both Humira and Enbrel unless one (1) of the exceptions on the PA form is present. For FDA-approved indications, an additional ninety (90) day trial of Cosentyx will also be required.		
ENBREL (etanercept)* CIMZIA (certolizumab pegol) *Full PA criteria may be found on the PA Criteria page by	ANTI-TNFs		
	ENBREL (etanercept)*	CIMZIA (certolizumab pegol)	*Full PA criteria may be found on the PA Criteria page by



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
HUMIRA (adalimumab)*	REMICADE (infliximab) RENFLEXIS (infliximab) SIMPONI subcutaneous (golimumab) OTHERS	clicking the hyperlink.
0005117/0//		*0
COSENTYX (secukinumab)	ACTEMRA subcutaneous (tocilizumab) ENTYVIO (vedolizumab) <sup>NR</sup> ILARIS (canakinumab) KEVZARA (sarilumab) KINERET (anakinra) ORENCIA subcutaneous (abatacept) OTEZLA (apremilast) SILIQ (brodalumab) STELARA subcutaneous (ustekinumab) TALTZ (ixekizumab) TREMFYA (guselkumab) XELJANZ (tofacitinib) XELJANZ XR (tofacitinib)	*Cosentyx will be authorized for treatment of plaque psoriasis, psoriatic arthritis and ankylosing spondylitis only after inadequate response to a ninety (90) day trial of Humira.
EPINEPHRINE, SELF-INJECTED	· ·	
<b>CLASS PA CRITERIA:</b> A non-preferred agent munderstand the training for the preferred agent(s).		tient's inability to follow the instructions, or the patient's failure to
epinephrine (labeler 49502 only)	ADRENACLICK (epinephrine) epinephrine (labeler 54505 and 00115) EPIPEN (epinephrine) EPIPEN JR (epinephrine)	
<b>ERYTHROPOIESIS STIMULATING</b>	* * * * * * * * * * * * * * * * * * * *	
<b>CLASS PA CRITERIA:</b> Non-preferred agents re PA form is present.	quire a thirty (30) day trial of a preferred agent before	re they will be approved, unless one (1) of the exceptions on the
EPOGEN (rHuEPO) PROCRIT (rHuEPO)	ARANESP (darbepoetin)	Erythropoiesis agents will be authorized if the following criteria are met:  1. Hemoglobin or Hematocrit less than 10/30 respectively. For renewal, hemoglobin or hematocrit levels greater than 12/36 will require dosage reduction or discontinuation. Exceptions will be considered on an individual basis after medical documentation is reviewed. (Lab oratory values must be dated within six (6) weeks of request.) and  2. Transferrin saturation ≥ 20%, ferritin levels ≥100 mg/ml, or on concurrent therapeutic iron therapy. (Laboratory values must be dated within three (3) weeks of request. For re-authorization, transferrin



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
		<ul> <li>saturation or ferritin levels are not required if the patient has been responsive to the erythropoietin agent and</li> <li>3. For HIV-infected patients, endogenous serum erythropoietin level must be ≤ 500mU/ml to initiate therapy and</li> <li>4. No evidence of untreated GI bleeding, hemolysis, or Vitamin B-12, iron or folate deficiency.</li> </ul>	
FLUOROQUINOLONES (Oral)AP			
<b>CLASS PA CRITERIA:</b> Non-preferred agents form is present.	require a five (5) day trial of a preferred agent before	they will be approved, unless one (1) of the exceptions on the PA	
CIPRO SUSPENSION (ciprofloxacin) ciprofloxacin levofloxacin tablet	AVELOX (moxifloxacin)  BAXDELA (delafloxacin)  CIPRO TABLETS (ciprofloxacin)  CIPRO XR (ciprofloxacin)  ciprofloxacin ER  ciprofloxacin suspension  FACTIVE (gemifloxacin)  LEVAQUIN (levofloxacin)  levofloxacin solution  moxifloxacin  NOROXIN (norfloxacin)  ofloxacin		
GLUCOCORTICOIDS, INHALEDAP	SHOXAGHT		
CLASS PA CRITERIA: Non-preferred agents exceptions on the PA form is present.	require thirty (30) day trials of each chemically unique	preferred agent before they will be approved, unless one (1) of the	
	GLUCOCORTICOIDS		
FLOVENT DISKUS (fluticasone) FLOVENT HFA (fluticasone) PULMICORT FLEXHALER (budesonide) PULMICORT RESPULES (budesonide)* QVAR REDIHALER (beclomethasone)	AEROSPAN (flunisolide)** ALVESCO (ciclesonide)  ARMONAIR RESPICLICK (fluticasone)  ARNUITY ELLIPTA (fluticasone)  ASMANEX HFA (mometasone)  ASMANEX TWISTHALER (mometasone)  budesonide	*Pulmicort Respules are only preferred for children up to nine (9) years of age. For patients nine (9) and older, prior authorization is required and will be approved only for a diagnosis of severe nasal polyps.  **Aerospan will be authorized for children ages 6 through 11 years old without a trial of a preferred agent.	
	GLUCOCORTICOID/BRONCHODILATOR COM	IRINATIONS	
ADVAIR DISKUS (fluticasone/salmeterol) ADVAIR HFA (fluticasone/salmeterol) DULERA (mometasone/formoterol)	AIRDUO RESPICLICK (fluticasone/salmeterol) BREO ELLIPTA (fluticasone/vilanterol) fluticasone/salmeterol	Substitute for Class Criteria: For a diagnosis of COPD only, non-preferred agents require sixty (60) day trials of each chemically unique preferred agent in this sub-class before they	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
SYMBICORT(budesonide/formoterol)		will be authorized, unless one (1) of the exceptions on the PA form is present. NOTE: Agents without an FDA-approved indication for COPD do not need to be trialed.
GROWTH HORMONE <sup>CL</sup>		
<b>CLASS PA CRITERIA:</b> Non-preferred agents red the PA form is present.	quire three (3) month trials of each preferred agent to	pefore they will be approved, unless one (1) of the exceptions on
GENOTROPIN (somatropin) NORDITROPIN (somatropin)	HUMATROPE (somatropin) INCRELEX (mecasermin) NUTROPIN AQ (somatropin) OMNITROPE (somatropin) SAIZEN (somatropin) SEROSTIM (somatropin) ZOMACTON (somatropin) ZORBTIVE (somatropin)	Patients already on a non-preferred agent will receive authorization to continue therapy on that agent for the duration of the existing PA.
H. PYLORI TREATMENT		
		components of the requested non-preferred agent and must be vill be approved, unless one (1) of the exceptions on the PA form
Please use individual components:     preferred PPI (omeprazole or pantoprazole)     amoxicillin     tetracycline     metronidazole     clarithromycin     bismuth	HELIDAC (bismuth/metronidazole/tetracycline) lansoprazole/amoxicillin/clarithromycin OMECLAMOX-PAK (omeprazole/amoxicillin/clarithromycin) PREVPAC (lansoprazole/amoxicillin/clarithromycin) PYLERA (bismuth/metronidazole/tetracycline)	
HEPATITIS B TREATMENTS		
	uire ninety (90) day trials of each preferred agent bef	ore they will be approved, unless one (1) of the exceptions on the
BARACLUDE (entecavir) lamivudine HBV	adefovir entecavir EPIVIR HBV (lamivudine) HEPSERA (adefovir) VEMLIDY (tenofovir alafenamide fumarate)	
HEPATITIS C TREATMENTS <sup>CL</sup>		
CLASS PA CRITERIA: For patients starting the require medical reasoning why a preferred regime		I on the PA Criteria page. Requests for non-preferred regimens
EPCLUSA (sofosbuvir/velpatasvir)* HARVONI (ledipasvir/sofosbuvir)* MAVYRET (pibrentasvir/glecaprevir)*	COPEGUS (ribavirin) DAKLINZA (daclatasvir)* MODERIBA 400 mg, 600 mg	*Full PA criteria may be found on the PA Criteria page by clicking the hyperlink.



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ribavirin ZEPATIER (elbasvir/grazoprevir)*	MODERIBA DOSE PACK PEGASYS (pegylated interferon) PEG-INTRON (pegylated interferon) OLYSIO (simeprevir)* REBETOL (ribavirin) RIBASPHERE RIBAPAK (ribavirin) RIBASPHERE 400 mg, 600 mg (ribavirin) SOVALDI (sofosbuvir)* TECHNIVIE (ombitasvir/paritaprevir/ritonavir)* VIEKIRA PAK (dasabuvir/ombitasvir/paritaprevir/ritonavir)* VIEKIRA XR (dasabuvir/ombitasvir/paritaprevir/ritonavir)* VIEKIRA XR (dasabuvir/ombitasvir/paritaprevir/ritonavir)*	
HYPERPARATHYROID AGENTSAP	,	
<b>CLASS PA CRITERIA:</b> Non-preferred agents re-PA form is present.	quire thirty (30) day trials of each preferred agent bef	fore they will be approved, unless one (1) of the exceptions on the
doxercalciferol paricalcitol capsule	HECTOROL (doxercalciferol) paricalcitol injection RAYALDEE (calcifediol) SENSIPAR (cinacalcet) ZEMPLAR (paricalcitol)	
HYPOGLYCEMICS, BIGUANIDES	ZZIM Zi w (panoaono)	
	quire a ninety (90) day trial of a preferred agent of si	imilar duration before they will be approved, unless one (1) of the
metformin ER (generic Glucophage XR)	FORTAMET (metformin ER) GLUCOPHAGE (metformin) GLUCOPHAGE XR (metformin ER) GLUMETZA (metformin ER)* metformin ER (generic Glumetza & Fortamet) RIOMET (metformin)	*Glumetza will be approved only after a 30-day trial of Fortamet.
<b>HYPOGLYCEMICS, DPP-4 INHIBIT</b>		
CLASS PA CRITERIA: Non-preferred agents a NOTE: DPP-4 inhibitors will NOT be approved	·	
JANUMET (sitagliptin/metformin) JANUVIA (sitagliptin) JENTADUETO (linagliptin/metformin) TRADJENTA (linagliptin)	alogliptin alogliptin/metformin alogliptin/pioglitazone JANUMET XR (sitagliptin/metformin)	



EFFECTIVE 07/01/2018 Version 2018.3e

This is not an all-inclusive list of available covered drugs and includes only managed categories. Refer to cover page for complete list of rules governing this PDL.

THERAPEUTIC DRUG CLASS			
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
	JENTADUETO XR (linagliptin/metformin) KAZANO (alogliptin/metformin) KOMBIGLYZE XR (saxagliptin/metformin) NESINA (alogliptin) ONGLYZA (saxagliptin) OSENI (alogliptin/pioglitazone)		

#### HYPOGLYCEMICS, GLP-1 AGONISTSCL

CLASS PA CRITERIA: Agents in this class will not be approved for patients with a starting A1C < 7%. Non-preferred agents are available only on appeal. Preferred agents in this class shall be approved in six (6) month intervals if the following criteria are met:

- Initial starts require a diagnosis of Type 2 Diabetes and an A1C taken within the last 30 days reflecting the patient's current and stabilized regimen. Current A1C must be ≤ 9%
- No agent in this class shall be approved except as add on therapy to a regimen consisting of at least one (1) other agent prescribed at the maximum tolerable dose for at least 90 days.
- Re-authorizations require <u>continued</u> maintenance on a regimen consisting of at least one (1) other agent at the maximum tolerable dose AND an A1C of ≤8%.

NOTE: GLP-1 agents will NOT be approved in combination with a DPP-4 inhibitor.

BYDUREON (exenatide)	ADLYXIN (lixisenatide)	
,	(	
BYETTA (exenatide)	TANZEUM (albiglutide)	
(	(3, 1, 3, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	
VICTOZA (liraglutide)	TRULICITY (dulaglutide)	
	(2222)	

### HYPOGLYCEMICS, INSULIN AND RELATED AGENTS

**CLASS PA CRITERIA:** Non-preferred agents require a ninety (90) day trial of a pharmacokinetically similar agent before they will be approved, unless one (1) of the exceptions on the PA form is present.

Humulin pens and Humalog Mix pens will be authorized only for patients who cannot utilize vials due to impaired vision or dexterity.				
HUMALOG (insulin lispro)	AFREZZA (insulin) <sup>CL</sup>	*Apidra will be authorized if the following criteria are met:		
HUMALOG MIX VIALS (insulin lispro/lispro	APIDRA (insulin glulisine) <sup>AP*</sup>	1. Patient is four (4) years of age or older; and		
protamine)	BASAGLAR (insulin glargine)	2. Patient is currently on a regimen including a longer		
HUMULIN VIALS (insulin)	FIASP (insulin aspart)	acting or basal insulin, and		
LANTUS (insulin glargine)	HUMALOG JR KWIKPEN (insulin lispro)	3. Patient has had a trial of a similar preferred agent,		
LEVEMIR (insulin detemir)	HUMALOG PEN/KWIKPEN (insulin lispro)	Novolog or Humalog, with documentation that the		
NOVOLOG (insulin aspart)	HUMALOG MIX PENS (insulin lispro/lispro	desired results were not achieved.		
NOVOLOG MIX (insulin aspart/aspart	protamine)			
protamine)	HUMULIN PENS (insulin)	**Tresiba U-100 will be authorized only for patients with a 6-		
	NOVOLIN (insulin)	month history of compliance on preferred long-acting insulin.		
	SOLIQUA (insulin glargine/lixisenatide)***			
	TOUJEO SOLOSTAR (insulin glargine)**	Tresiba U-200 and Toujeo Solostar will <b>only</b> be approved for		
	TRESIBA (insulin degludec)**	patients with a 6-month history of compliance on preferred		
	XULTOPHY (insulin degludec/liraglutide)***	long-acting insulin who require once-daily doses of at least		



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS				
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA		
		60 units of insulin.		
		***Non-preferred insulin combination products require that the patient must already be established on the individual agents at doses not exceeding the maximum dose achievable with the combination product, and require medical reasoning beyond convenience or enhanced compliance as to why the clinical need cannot be met with a combination of preferred single-ingredient agents.		
HYPOGLYCEMICS, MEGLITINIDES				
CLASS PA CRITERIA: Non-preferred agents ar				
nateglinide	MEGLITINIDES PRANDIN (repaglinide)			
repaglinide	STARLIX (nateglinide)			
	MEGLITINIDE COMBINATIONS			
	PRANDIMET (repaglinide/metformin) repaglinide/metformin			
HYPOGLYCEMICS, MISCELLANEO	US AGENTS			
<b>CLASS PA CRITERIA:</b> Welchol will be authorized agent.	d for add-on therapy for type 2 diabetes when there is	s a previous history of a thirty (30) day trial of an oral diabetic		
WELCHOL (colesevelam) <sup>AP</sup>	SYMLIN (pramlintide)*	*Symlin will be authorized with a history of bolus insulin utilization in the past ninety (90) days with no gaps in insulin therapy greater than thirty (30) days.		
HYPOGLYCEMICS, SGLT2 INHIBITO	ORS <sup>CL</sup>			
CLASS PA CRITERIA: Agents in this class will not be approved for patients with a starting A1C < 7%. Non-preferred agents are available only on appeal. Preferred agents in this class shall be approved in six (6) month intervals if the following criteria are met.				
<ul> <li>Initial starts require a diagnosis of Type 2 Diabetes and an A1C taken within the last 30 days reflecting the patient's current and stabilized regimen. Current A1C must be ≤ 9%.</li> </ul>				
<ul> <li>No agent in this class shall be approved exce dose for at least 90 days.</li> </ul>	ept as add on therapy to a regimen consisting of at le	ast one (1) other agent prescribed at the maximum tolerable		
• Re-authorizations require <u>continued</u> maintenance on a regimen consisting of at least one (1) other agent at the maximum tolerable dose AND an A1C of ≤8%.				
SGLT2 INHIBITORS				
FARXIGA (dapagliflozin) JARDIANCE (empagliflozin)	INVOKANA (canagliflozin)			



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	SGLT2 COMBINATIONS	
SYNJARDY (empagliflozin/metformin) SYNJARDY XR (empagliflozin/metformin)	GLYXAMBI (empagliflozin/linagliptin) INVOKAMET (canagliflozin/metformin) INVOKAMET XR (canagliflozin/metformin) XIGDUO XR (dapagliflozin/metformin)	
HYPOGLYCEMICS, TZD	,	
CLASS PA CRITERIA: Non-preferred agents a	re available only on appeal.	
	THIAZOLIDINEDIONES	
pioglitazone	ACTOS (pioglitazone) AVANDIA (rosiglitazone)	
	TZD COMBINATIONS	
	ACTOPLUS MET (pioglitazone/ metformin) ACTOPLUS MET XR (pioglitazone/ metformin) AVANDAMET (rosiglitazone/metformin) AVANDARYL (rosiglitazone/glimepiride) DUETACT (pioglitazone/glimepiride) pioglitazone/glimepiride pioglitazone/ metformin	Patients are required to use the components of Actoplus Met and Duetact separately. Exceptions will be handled on a case-by-case basis.
IMMUNOMODULATORS, ATOPIC D	• •	
CLASS PA CRITERIA: Non-preferred agents re	quire 30-day trial of a medium to high potency topica	al corticosteroid <b>AND all</b> preferred agents in this class unless one ded with involvement of sensitive areas such as the face and skin
ELIDEL (pimecrolimus) EUCRISA (crisaborole) <sup>AP*</sup>	DUPIXENT (dupilumab)** PROTOPIC (tacrolimus)*** tacrolimus ointment	*Eucrisa requires a 30-day trial of Elidel <b>OR</b> a medium to high potency corticosteroid unless contraindicated.
	tacionina cinamoni	**Full PA criteria for Dupixent may be found on the PA Criteria page by clicking the hyperlink
		***Protopic brand is preferred over its generic equiviliant.
IMMUNOMODULATORS, GENITAL	WARTS & ACTINIC KERATOSIS AGE	NTS
<b>CLASS PA CRITERIA:</b> Non-preferred agents red PA form is present.	quire thirty (30) day trials of each preferred agent before	ore they will be approved, unless one (1) of the exceptions on the
CONDYLOX GEL (podofilox) EFUDEX (fluorouracil) imiquimod	ALDARA (imiquimod) CARAC (fluorouracil) diclofenac 3% gel fluorouracil 0.5% cream fluorouracil 5% cream podofilox SOLARAZE (diclofenac) TOLAK (fluorouracil 4% cream)	*Zyclara will be authorized for a diagnosis of actinic keratosis.



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS				
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA		
	VEREGEN (sinecatechins) ZYCLARA (imiquimod)*			
IMMUNOSUPPRESSIVES, ORAL				
CLASS PA CRITERIA: Non-preferred agents require a fourteen (14) day trial of a preferred agent before they will be approved, unless one (1) of the exceptions on the PA form is present.				
azathioprine cyclosporine cyclosporine, modified mycophenolate mofetil sirolimus tacrolimus capsule	ASTAGRAF XL (tacrolimus) AZASAN (azathioprine) CELLCEPT (mycophenolate mofetil) ENVARSUS XR (tacrolimus) IMURAN (azathioprine) mycophenolic acid mycophenolic mofetil suspension MYFORTIC (mycophenolic acid) NEORAL (cyclosporine, modified) PROGRAF (tacrolimus) RAPAMUNE (sirolimus) SANDIMMUNE (cyclosporine) ZORTRESS (everolimus)			
INTRANASAL RHINITIS AGENTS <sup>AP</sup>				
CLASS PA CRITERIA: See below for individual	sub-class criteria.			
	ANTICHOLINERGICS			
ipratropium	ATROVENT(ipratropium)	Non-preferred agents require thirty (30) day trials of one (1) preferred nasal anti-cholinergic agent, <b>AND</b> one (1) preferred antihistamine <b>AND</b> one (1) preferred intranasal corticosteroid agent before they will be approved, unless one (1) of the exceptions on the PA form is present.		
	ANTIHISTAMINES			
azelastine	ASTEPRO (azelastine) olopatadine PATANASE (olopatadine)	Non-preferred agents require thirty (30) day trials of one (1) preferred antihistamine <b>AND</b> one (1) preferred intranasal corticosteroid before they will be approved, unless one (1) of the exceptions on the PA form is present.		
COMBINATIONS				
	DYMISTA (azelastine / fluticasone)	Dymista requires a concurrent thirty (30) day trial of each preferred component before it will be approved, unless one (1) of the exceptions on the PA form is present.		
CORTICOSTEROIDS				
fluticasone propionate	BECONASE AQ (beclomethasone) budesonide FLONASE (fluticasone propionate) flunisolide	Non-preferred agents require thirty (30) day trials of the preferred agent in this sub-class before they will be approved, unless one (1) of the exceptions on the PA form is present		



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	S
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	mometasone NASACORT AQ (triamcinolone) NASONEX (mometasone) OMNARIS (ciclesonide) QNASL HFA (beclomethasone) RHINOCORT AQUA (budesonide) triamcinolone VERAMYST (fluticasone furoate) ZETONNA (ciclesonide)	
IRRITABLE BOWEL SYNDROME/S	HORT BOWEL SYNDROME/SELECTE	D GI AGENTS CL
CLASS PA CRITERIA: All agents are approvable	e only for patients age eighteen (18) and older. See b	elow for additional sub-class criteria.
	CONSTIPATION	
AMITIZA (lubiprostone)* MOVANTIK (naloxegol)**	LINZESS (linaclotide)*** RELISTOR INJECTION (methylnaltrexone)**** RELISTOR TABLET (methylnaltrexone)**** SYMPROIC (naldemedine) TRULANCE (plecanatide)*****	All agents require documentation of the current diagnosis and evidence that the patient has failed to find relief with dietary modification and a fourteen (14) day trial of an osmotic laxative.  In addition:  * Amitiza is indicated for CIC, IBS-C (females only) and OIC. Approval for the diagnosis of OIC requires a concurrent and continuous 90-day history of opioid claims on record.  ** Movantik will be approved per the FDA-approved label for OIC with a concurrent and continuous 90-day history of opioid claims on record.  *** Linzess is indicated for CIC and IBS-C and requires a thirty (30) day trial of Amitiza For the indication of IBS-C in males, a trial of Amitiza is not requires thirty (30) day trials of both Movantik and Amitiza.  ***** Trulance is indicated for CIC and requires a thirty (30) day trial of Amitiza. For the indication of IBS-C in males, a trial of Amitiza is not required.
	DIARRHEA	
	alosetron* MYTESI (crofelemer)* LOTRONEX (alosetron)* VIBERZI (eluxadoline)*	* Full PA criteria may be found on the <u>PA Criteria</u> page by clicking the hyperlink.



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
LAXATIVES AND CATHARTICS		
CLASS PA CRITERIA: Non-preferred agents red PA form is present	quire thirty (30) day trials of each preferred agent before	ore they will be approved, unless one (1) of the exceptions on the
COLYTE GOLYTELY NULYTELY peg 3350	HALFLYTELY-BISACODYL KIT MOVIPREP OSMOPREP PREPOPIK SUPREP	
LEUKOTRIENE MODIFIERS		
<b>CLASS PA CRITERIA:</b> Non-preferred agents red PA form is present.	quire thirty (30) day trials of each preferred agent before	ore they will be approved, unless one (1) of the exceptions on the
montelukast zafirlukast	ACCOLATE (zafirlukast) SINGULAIR (montelukast) zileuton ZYFLO (zileuton)	
LIPOTROPICS, OTHER (Non-statin		
<b>CLASS PA CRITERIA:</b> Non-preferred agents red PA form is present.	quire a twelve (12) week trial of a preferred agent before	ore they will be approved, unless one (1) of the exceptions on the
·	BILE ACID SEQUESTRANTSAP	
cholestyramine colestipol tablets	COLESTID (colestipol) colestipol granules KYNAMRO (mipomersen)* QUESTRAN (cholestyramine) WELCHOL (colesevelam)**	*Full PA criteria may be found on the PA Criteria page by clicking the hyperlink.  **Welchol will be authorized for add-on therapy for type 2 diabetes when there is a previous history of a thirty (30) day trial of an oral agent (metformin, sulfonylurea or thiazolidinedione (TZD)). See HYPOGLYCEMICS, MISCELLANEOUS.
	CHOLESTEROL ABSORPTION INHIBIT	
ZETIA (ezetimibe) AP	ezetimibe	Zetia will be authorized with prior use of a HMG-CoA reductase inhibitor within the previous six (6) months.
FATTY ACIDS <sup>AP</sup>		
	LOVAZA (omega-3-acid ethyl esters) omega-3 acid ethyl esters VASCEPA (icosapent ethyl)	These agents shall only be authorized when the patient has an initial triglyceride level ≥ 500 mg/dL and has had inadequate response or intolerance to trials of BOTH a nicotinic acid and a fibrate, unless otherwise contraindicated.
formatile and a 40 mm	FIBRIC ACID DERIVATIVES <sup>AP</sup>	
fenofibrate 40 mg fenofibrate 54, 150 and 160 mg fenofibrate micronized 67mg, 134mg & 200mg	ANTARA (fenofibrate) FENOGLIDE (fenofibrate) FIBRICOR (fenofibric acid)	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
fenofibrate nanocrystallized 48 mg, 145 mg gemfibrozil	fenofibrate 43, 50, 120 and 130 mg fenofibric acid LIPOFEN (fenofibrate) LOFIBRA (fenofibrate) LOPID (gemfibrozil) TRICOR (fenofibrate nanocrystallized) TRIGLIDE (fenofibrate) TRILIPIX (fenofibric acid)	
	MTP INHIBITORS  JUXTAPID (lomitapide)*	*Full PA criteria may be found on the PA Criteria page by
	OOXITA ID (IoIIIIapide)	clicking the hyperlink.
	NIACIN	
niacin NIACOR (niacin) NIASPAN (niacin)	niacin ER	
	PCSK-9 INHIBITORS	
	PRALUENT (alirocumab)* REPATHA (evolocumab)*	*Full PA criteria may be found on the <u>PA Criteria</u> page by clicking the hyperlink.
LIPOTROPICS, STATINS <sup>AP</sup>		
CLASS PA CRITERIA: See below for individual s	sub-class criteria.	
	STATINS	
atorvastatin lovastatin pravastatin rosuvastatin simvastatin*	ALTOPREV (lovastatin) CRESTOR (rosuvastatin) fluvastatin fluvastatin ER LESCOL (fluvastatin) LESCOL XL (fluvastatin) LIPITOR (atorvastatin) LIVALO (pitavastatin) MEVACOR (lovastatin) PRAVACHOL (pravastatin) ZOCOR (simvastatin)*	Non-preferred agents require twelve (12) week trials of two (2) preferred agents, including the generic formulation of the requested non-preferred agent, before they will be approved, unless one (1) of the exceptions on the PA form is present.  *Zocor/simvastatin 80mg tablets will require a clinical PA.
STATIN COMBINATIONS		
	ADVICOR (lovastatin/niacin) amlodipine/atorvastatin CADUET (atorvastatin/amlodipine) ezetimibe/simvastatin LIPTRUZET (atorvastatin/ezetimibe) SIMCOR (simvastatin/niacin ER)	Non-preferred agents require thirty (30) day concurrent trials of the corresponding preferred single agents before they will be approved, unless one (1) of the exceptions on the PA form is present.  *Vytorin will be authorized only after an insufficient response to
		*Vytorin will be authorized only after an insufficient response a twelve (12) week trial of the maximum tolerable dose



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
		atorvastatin or rosuvastatin, unless one (1) of the exceptions on the PA form is present.
		Vytorin 80/10mg tablets will require a clinical PA.
	s require a five (5) day trial of each preferred agent be	fore they will be approved, unless one (1) of the exceptions on the
PA form is present.	MACROLIDES	
azithromycin clarithromycin suspension erythromycin base	BIAXIN (clarithromycin) clarithromycin tablets clarithromycin ER E.E.S. (erythromycin ethylsuccinate) E-MYCIN (erythromycin) ERYC (erythromycin) ERYPED (erythromycin ethylsuccinate) ERY-TAB (erythromycin) ERYTHROCIN (erythromycin stearate) erythromycin estolate PCE (erythromycin) ZITHROMAX (azithromycin) ZMAX (azithromycin)	
<b>MULTIPLE SCLEROSIS AGENTS</b>		
	s require a diagnosis of multiple sclerosis and thirty (30) s one (1) of the exceptions on the PA form is present.	day trials of each chemically unique preferred agent in the same
	INTERFERONS <sup>AP</sup>	
AVONEX (interferon beta-1a) AVONEX PEN (interferon beta-1a) BETASERON (interferon beta-1b)	EXTAVIA KIT (interferon beta-1b) EXTAVIA VIAL (interferon beta-1b) PLEGRIDY (peginterferon beta-1a) REBIF (interferon beta-1a) REBIF REBIDOSE (interferon beta-1a)	
NON-INTERFERONS		
COPAXONE 20 mg (glatiramer) GILENYA (fingolimod) *	AMPYRA (dalfampridine)** AUBAGIO (teriflunomide)*** COPAXONE 40 mg (glatiramer)**** GLATOPA (glatiramer) TECFIDERA (dimethyl fumarate)**** ZINBRYTA (daclizumab)	In addition to class PA criteria, the following conditions and criteria also apply:  *Gilenya will be approved after a thirty (30) day trial of a preferred injectable agent.  **Ampyra will be authorized if the following criteria are met:  1. Diagnosis of multiple sclerosis and  2. No history of seizures and  3. No evidence of moderate or severe renal impairment



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
PREFERRED AGENTS	NON-PREFERRED AGENTS	and 4. Initial prescription will be authorized for thirty (30) days only.  ***Aubagio will be authorized if the following criteria are met:  1. Diagnosis of relapsing multiple sclerosis and  2. Measurement of transaminase and bilirubin levels within the (6) months before initiation of therapy and ALT levels at least monthly for six (6) months after initiation of therapy and  3. Complete blood cell count (CBC) within six (6) months before initiation of therapy and  4. Female patients must have a negative pregnancy test before initiation of therapy and be established on a reliable method of contraception if appropriate and  5. Patient is from eighteen (18) up to sixty-five (65) years of age and  6. Negative tuberculin skin test before initiation of therapy  ****Copaxone 40mg will only be authorized for documented injection site issues.  *****Tecfidera will be authorized if the following criteria are met:
		<ol> <li>Diagnosis of relapsing multiple sclerosis and</li> <li>Complete blood count (CBC) within six (6) months of initiation of therapy and six (6) months after initiation and</li> </ol>
NEUROPATHIC PAIN		<ol><li>Complete blood count (CBC) annually during therapy.</li></ol>
	uire a trial of a preferred agent in the corresponding	dosage form (oral or topical) before they will be approved, unless
capsaicin OTC duloxetine gabapentin lidocaine patch	CYMBALTA (duloxetine) GRALISE (gabapentin)* HORIZANT (gabapentin) IRENKA (duloxetine) LIDODERM (lidocaine) LYRICA CAPSULE (pregabalin)** LYRICA SOLUTION (pregabalin)** NEURONTIN (gabapentin) QUTENZA (capsaicin) SAVELLA (milnacipran)***	*Gralise will be authorized if the following criteria are met:  1. Diagnosis of post herpetic neuralgia and 2. Trial of a tricyclic antidepressant for a least thirty (30) days and 3. Trial of gabapentin immediate release formulation (positive response without adequate duration) and 4. Request is for once daily dosing with 1800 mg maximum daily dosage.  **Lyrica will be authorized if the following criteria are met:



EFFECTIVE 07/01/2018 Version 2018.3e

This is not an all-inclusive list of available covered drugs and includes only managed categories. Refer to cover page for complete list of rules governing this PDL.

THERAPEUTIC DRUG CLASS

	THERAI EO NO DROG GEAC	
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	ZOSTRIX OTC (capsaicin)	1. Diagnosis of seizure disorders or neuropathic pain associated with a spinal cord injury or  2. Diagnosis of fibromyalgia, postherpetic neuralgia, or diabetic neuropathy AND a history of a trial of duloxetine at the generally accepted maximum therapeutic dose of 60 mg/day AND gabapentin at a therapeutic dose range between 900 mg and 2,400 mg per day for thirty (30) days within the previous twenty-four (24) month period or an intolerance due to a potential adverse drug-drug interaction, drug-disease interaction, or intolerable side effect (In cases of renal impairment, doses may be adjusted based on the degree of impairment.)  ***Savella will be authorized for a diagnosis of fibromyalgia or a previous thirty (30) day trial of a drug that infers fibromyalgia:
NSAIDS <sup>AP</sup>		duloxetine, gabapentin, amitriptyline or nortriptyline.
CLASS PA CRITERIA: See below for sub-class	PA criteria.	
	NON-SELECTIVE	
diclofenac (IR, SR) flurbiprofen ibuprofen (Rx and OTC) INDOCIN SUSPENSION (indomethacin) indomethacin ketoprofen ketorolac meloxicam tablet nabumetone naproxen (Rx and OTC) piroxicam sulindac	ANAPROX (naproxen) ANSAID (flurbiprofen) CATAFLAM (diclofenac) CLINORIL (sulindac) DAYPRO (oxaprozin) diflunisal DUEXIS (famotidine/ibuprofen) etodolac IR etodolac SR FELDENE (piroxicam) fenoprofen INDOCIN SUPPOSITORIES (indomethacin) indomethacin ER ketoprofen ER LODINE (etodolac) meclofenamate mefenamic acid meloxicam suspension MOBIC TABLET (meloxicam) MOTRIN (ibuprofen) NALFON (fenoprofen)	Non-preferred agents require thirty (30) day trials of each preferred agent before they will be approved, unless one (1) of the exceptions on the PA form is present.



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	SS
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	NAPRELAN (naproxen) NAPROSYN (naproxen) naproxen CR oxaprozin PONSTEL (meclofenamate) SPRIX (ketorolac) TIVORBEX (indomethacin) tolmetin VIVLODEX (meloxicam) VOLTAREN (diclofenac) ZIPSOR (diclofenac potassium) ZORVOLEX (diclofenac)	
	NSAID/GI PROTECTANT COMBINATIO	
	ARTHROTEC (diclofenac/misoprostol) diclofenac/misoprostol VIMOVO (naproxen/esomeprazole)	Non-preferred agents are only available on appeal and require medical reasoning beyond convenience as to why the need cannot be met with the combination of preferred single agents.
	COX-II SELECTIVE	
	CELEBREX (celecoxib) celecoxib	COX-II Selective agents require thirty (30) day trials of each preferred Non-Selective Oral NSAID, <b>UNLESS</b> the following criteria are met:  Patient has a history or risk of a serious GI complication; <b>OR</b> Agent is requested for treatment of a chronic condition <b>and</b> 1. Patient is seventy (70) years of age or older, <b>or</b> 2. Patient is currently on anticoagulation therapy.
	TOPICAL	
VOLTAREN GEL (diclofenac)*	diclofenac gel diclofenac solution FLECTOR PATCH (diclofenac)** PENNSAID (diclofenac)	*Voltaren Gel will be limited to 100 grams per month.  Non-preferred agents require a thirty (30) day trial of the preferred Topical agent and thirty (30) day trials of each preferred oral NSAID before they will be approved, unless one (1) of the exceptions on the PA form is present.  **Flector patches will only be authorized for a diagnosis of acute strain, sprain or injury after a five (5) day trial of one (1) of the preferred oral NSAIDs and for a maximum duration of fourteen (14) days unless one (1) of the exceptions on the PA form is present.



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
OPHTHALMIC ANTIBIOTICSAP		
<b>CLASS PA CRITERIA:</b> Non-preferred agents re PA form is present.	quire three (3) day trials of each preferred agent before	ore they will be approved, unless one (1) of the exceptions on the
bacitracin/polymyxin ointment ciprofloxacin* erythromycin gentamicin levofloxacin* neomycin/bacitracin/polymyxin ofloxacin* polymyxin/trimethoprim sulfacetamide drops tobramycin TOBREX OINT (tobramycin)	AZASITE (azithromycin) bacitracin BLEPH-10 (sulfacetamide) BESIVANCE (besifloxacin)* CILOXAN (ciprofloxacin) GARAMYCIN (gentamicin) gatifloxacin ILOTYCIN (erythromycin) MOXEZA (moxifloxacin)** moxifloxacin** NATACYN (natamycin) neomycin/polymyxin/gramicidin NEOSPORIN (neomycin/polymyxin/gramicidin) OCUFLOX (ofloxacin) POLYTRIM (polymyxin/trimethoprim) sulfacetamide ointment TOBREX (tobramycin) VIGAMOX (moxifloxacin) ZYMAXID (gatifloxacin)	*Prior authorization of any fluoroquinolone agent requires three (3) day trials of all other preferred agents unless definitive laboratory cultures exist indicating the need to use a fluoroquinolone.  **Brand Vigamox will be preferred over Brand Moxeza, and both brands are preferred over their generic equivalent.
<b>OPHTHALMIC ANTIBIOTIC/STERO</b>	ID COMBINATIONS <sup>AP</sup>	
<b>CLASS PA CRITERIA:</b> Non-preferred agents re PA form is present.	quire three (3) day trials of each preferred agent before	ore they will be approved, unless one (1) of the exceptions on the
BLEPHAMIDE (prednisolone/sulfacetamide) neomycin/polymyxin/dexamethasone sulfacetamide/prednisolone TOBRADEX OINTMENT (tobramycin/ dexamethasone) TOBRADEX SUSPENSION (tobramycin/ dexamethasone)	BLEPHAMIDE S.O.P. (prednisolone/sulfacetamide) MAXITROL ointment (neomycin/polymyxin/dexamethasone) MAXITROL suspension (neomycin/polymyxin/dexamethasone) neomycin/bacitracin/polymyxin/hydrocortisone neomycin/polymyxin/hydrocortisone PRED-G (prednisolone/gentamicin) TOBRADEX ST (tobramycin/dexamethasone) tobramycin/dexamethasone suspension ZYLET (loteprednol/tobramycin)	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
OPHTHALMICS FOR ALLERGIC CO	ONJUNCTIVITIS <sup>AP</sup>	
<b>CLASS PA CRITERIA:</b> Non-preferred agents re of the exceptions on the PA form is present.	quire thirty (30) day trials of three (3) preferred chem	nically unique agents before they will be approved, unless one (1)
ALAWAY (ketotifen) cromolyn ketotifen olopatadine (Sandoz brand labeler 61314) ZADITOR OTC (ketotifen)	ALAMAST (pemirolast) ALOCRIL (nedocromil) ALOMIDE (lodoxamide) ALREX (loteprednol) azelastine BEPREVE (bepotastine) CROLOM (cromolyn) ELESTAT (epinastine) EMADINE (emedastine) epinastine LASTACAFT (alcaftadine) olopatadine (all labelers except Sandoz) OPTICROM (cromolyn) OPTIVAR (azelastine) PATADAY (olopatadine) PATANOL (olopatadine) PAZEO (olopatadine)	
OPHTHALMICS, ANTI-INFLAMMATORIES- IMMUNOMODULATORS		
CLASS PA CRITERIA: See below for individual		
	RESTASIS (cyclosporine) XIIDRA (lifitegrast)	<ol> <li>The following prior authorization criteria apply to both Restasis and Xiidra:</li> <li>Patient must be sixteen (16) years of age or greater; AND</li> <li>Prior Authorization must be requested by an ophthalmologist or optometrist; AND</li> <li>Clinically diagnosed tear deficiency due to ocular inflammation in patients with keratoconjunctivitis sicca or dry eye syndrome (also known as dry eye); AND</li> <li>Patient must have a functioning lacrimal gland; AND</li> <li>Patient using artificial tears at least four (4) times a day over the last thirty (30) days; AND</li> <li>Patient must not have an active ocular infection</li> </ol>



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS			
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
OPHTHALMICS, ANTI-INFLAMMAT	ORIES		
	equire five (5) day trials of at least two (2) preferrinclude at least one agent with the same mechanism	red agents before they will be approved, unless one (1) of the of action as the requested non-preferred agent.	
dexamethasone diclofenac DUREZOL (difluprednate) fluorometholone flurbiprofen ketorolac prednisolone acetate prednisolone sodium phosphate	ACULAR (ketorolac) ACULAR LS (ketorolac) ACUVAIL (ketorolac tromethamine) BROMDAY (bromfenac) bromfenac BROMSITE (bromfenac) FLAREX (fluorometholone) FML (fluorometholone) FML FORTE (fluorometholone) ILEVRO (nepafenac) LOTEMAX DROPS, OINTMENT (loteprednol) LOTEMAX GEL (loteprednol) MAXIDEX (dexamethasone) NEVANAC (nepafenac) OMNIPRED (prednisolone) OZURDEX (dexamethasone) PRED FORTE (prednisolone) PRED MILD (prednisolone) PROLENSA (bromfenac) RETISERT (fluocinolone) TRIESENCE (triamcinolone) VEXOL (rimexolone) XIBROM (bromfenac)		
OPHTHALMICS, GLAUCOMA AGEI	NTS		
CLASS PA CRITERIA: Non-preferred agents will only be authorized if there is an allergy to all preferred agents in the corresponding sub-class.			
COMPLOAN /haire anidir - //	COSORT (described friends)		
COMBIGAN (brimonidine/timolol) dorzolamide/timolol SIMBRINZA (brinzolamide/brimonidine)	COSOPT (dorzolamide/timolol) COSOPT PF (dorzolamide/timolol)		
DETORTIO O (L. L. L.)	BETA BLOCKERS		
BETOPTIC S (betaxolol) carteolol levobunolol timolol drops	BETAGAN (levobunolol) betaxolol BETIMOL (timolol) ISTALOL (timolol) OPTIPRANOLOL (metipranolol)		



EFFECTIVE 07/01/2018 Version 2018.3e

	THERAPEUTIC DRUG CLAS	SS
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	timolol gel TIMOPTIC (timolol)	
	CARBONIC ANHYDRASE INHIBITOR	RS
AZOPT (brinzolamide) orzolamide	TRUSOPT (dorzolamide)	
	PARASYMPATHOMIMETICS	
PHOSPHOLINE IODIDE (echothiophate iodide)	pilocarpine	
	PROSTAGLANDIN ANALOGS	
latanoprost TRAVATAN-Z (travoprost)	bimatoprost LUMIGAN (bimatoprost) RESCULA (unoprostone) travoprost XALATAN (latanoprost) ZIOPTAN (tafluprost)	
	SYMPATHOMIMETICS	
brimonidine 0.2%	ALPHAGAN P 0.1% Solution (brimonidine) ALPHAGAN P 0.15% Solution (brimonidine) apraclonidine brimonidine 0.15% IOPIDINE (apraclonidine)	
OPIATE DEPENDENCE TREATMEN CLASS PA CRITERIA: Buprenorphine/naloxone		with a documented intolerance of or allergy to Suboxone strips.
WV Medicaid's buprenorphine coverage policy ma	ay be viewed by clicking on the following hyperlink: B	uprenorphine Coverage Policy and Related Forms
Naloxone NARCAN NASAL SPRAY (naloxone) SUBOXONE FILM (buprenorphine/naloxone)*	buprenorphine tablets buprenorphine/naloxone tablets BUNAVAIL (buprenorphine/naloxone)	* Full PA criteria may be found on the PA Criteria page by clicking the hyperlink.
VIVITROL (naltrexone)	SUBLOCADE (buprenorphine soln)*NR ZUBSOLV (buprenorphine/naloxone)	VIVITROL no longer requires a PA.
OTIC ANTIBIOTICSAP		
<b>CLASS PA CRITERIA:</b> Non-preferred agents re PA form is present.	quire five (5) day trials of each preferred agent befo	re they will be approved, unless one (1) of the exceptions on the
CIPRO HC (ciprofloxacin/hydrocortisone) CIPRODEX (ciprofloxacin/dexamethasone) COLY-MYCIN S (colistin/hydrocortisone/ neomycin/thonzonium bromide) ofloxacin	ciprofloxacin CORTISPORIN-TC (colistin/hydrocortisone/ neomycin) neomycin/polymyxin/HC solution/suspension OTIPRIO VIAL (ciprofloxacin) OTOVEL (ciprofloxacin/fluocinolone)	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
<b>PAH AGENTS – ENDOTHELIN REC</b>	EPTOR ANTAGONISTSCL	
<b>CLASS PA CRITERIA:</b> Non-preferred agents red PA form is present.	quire a thirty (30) day trial of a preferred agent before	re they will be approved, unless one (1) of the exceptions on the
LETAIRIS (ambrisentan) TRACLEER (bosentan)	OPSUMIT (macitentan)	
PAH AGENTS – GUANYLATE CYCL	ASE STIMULATORCL	
CLASS PA CRITERIA: Non-preferred agents red of the exceptions on the PA form is present.	uire a thirty (30) day trial of a preferred agent from a	any other PAH Class before they will be approved, unless one (1)
	ADEMPAS (riociguat)	
PAH AGENTS – PDE5scl		
CLASS PA CRITERIA: Non-preferred agents red PA form is present. Patients stabilized on non-preferred agents will be		re they will be approved, unless one (1) of the exceptions on the
sildenafil	ADCIRCA (tadalafil) REVATIO IV (sildenafil) REVATIO SUSPENSION (sildenafil) REVATIO TABLETS (sildenafil)	
PAH AGENTS - PROSTACYCLINSC	L	
CLASS PA CRITERIA: Non-preferred agents re available), before they will be approved, unless on		cluding the preferred generic form of the non-preferred agent (if
epoprostenol VENTAVIS (iloprost)*	FLOLAN (epoprostenol) ORENITRAM ER (treprostinil) REMODULIN (treprostinil sodium) TYVASO (treprostinil) UPTRAVI (selexipag) VELETRI (epoprostenol)	*Ventavis will only be authorized for the treatment of pulmonary artery hypertension (WHO Group 1) in patients with NYHA Class III or IV symptoms.
PANCREATIC ENZYMESAP		
<b>CLASS PA CRITERIA:</b> Non-preferred agents red PA form is present. For members with cystic fibrosis, a trial of a prefer		re they will be approved, unless one (1) of the exceptions on the
CREON ZENPEP	PANCREAZE PERTZYE ULTRESA VIOKACE	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
PHOSPHATE BINDERSAP		
CLASS PA CRITERIA: Non-preferred agents re exceptions on the PA form is present.	equire a thirty (30) day trial of at least two (2) prefe	rred agents before they will be approved, unless one (1) of the
calcium acetate MAGNEBIND RX (calcium carbonate, folic acid, magnesium carbonate) PHOSLYRA (calcium acetate) RENAGEL (sevelamer)	AURYXIA (ferric citrate) ELIPHOS (calcium acetate) FOSRENOL (lanthanum) PHOSLO (calcium acetate) RENVELA (sevelamer carbonate) sevelamer carbonate VELPHORO (sucroferric oxyhydroxide)	
PLATELET AGGREGATION INHIBIT	TORS	
<b>CLASS PA CRITERIA:</b> Non-preferred agents red PA form is present.	quire a thirty (30) day trial of a preferred agent before	re they will be approved, unless one (1) of the exceptions on the
AGGRENOX (dipyridamole/ASA) BRILINTA (ticagrelor) clopidogrel prasugrel	clopidogrel kit dipyridamole dipyridamole/aspirin DURLAZA ER (aspirin) EFFIENT (prasugrel) PERSANTINE (dipyridamole) PLAVIX (clopidogrel) TICLID (ticlopidine) ticlopidine ZONTIVITY (vorapaxar)	
PROGESTINS FOR CACHEXIA		
CLASS PA CRITERIA: Non-preferred agents require a thirty (30) day trial of a preferred agent before they will be approved, unless one (1) of the exceptions on the PA form is present.		
megestrol	MEGACE ES (megestrol)	
PROGESTATIONAL AGENTS		
	und on the PA Criteria page by clicking the hyperlink	•
MAKENA (hydroxyprogesterone caproate)		
PROTON PUMP INHIBITORS <sup>AP</sup>		
CLASS PA CRITERIA: Non-preferred agents require sixty (60) day trials of both omeprazole (Rx) and pantoprazole at the maximum recommended dose*, inclusive of a concurrent thirty (30) day trial at the maximum dose of an H <sub>2</sub> antagonist before they will be approved, unless one (1) of the exceptions on the PA form is present.		
omeprazole (Rx) pantoprazole	ACIPHEX (rabeprazole) ACIPHEX SPRINKLE (rabeprazole)	*Maximum recommended doses of the PPIs and H2-receptor antagonists may be located at the BMS Pharmacy PA criteria



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
PREVACID SOLUTABS (lansoprazole)** PROTONIX GRANULES (pantoprazole)	DEXILANT (dexlansoprazole) esomeprazole magnesium esomeprazole strontium lansoprazole Rx NEXIUM (esomeprazole) omeprazole/sodium bicarbonate (Rx) PREVACID CAPSULES (lansoprazole) PRILOSEC Rx (omeprazole) PROTONIX DR TABLETS (pantoprazole) rabeprazole ZEGERID Rx (omeprazole/sodium bicarbonate)	page titled "Max PPI and H2RA" by clicking on the hyperlink.  **Prior authorization is required for Prevacid Solutabs for members nine (9) years of age or older.
SEDATIVE HYPNOTICSAP		
	uire thirty (30) day trials of the preferred agent in <b>BO</b> ats in this class will be limited to fifteen (15) tablets in	<b>PTH</b> sub-classes before they will be approved, unless one (1) of a thirty (30) day period.
	BENZODIAZEPINES	
temazepam 15, 30 mg	DALMANE (flurazepam) DORAL (quazepam) estazolam flurazepam HALCION (triazolam) quazepam RESTORIL (temazepam) temazepam 7.5, 22.5 mg triazolam	
	OTHERS	
zolpidem 5, 10 mg	AMBIEN (zolpidem) AMBIEN CR (zolpidem) BELSOMRA (suvorexant) chloral hydrate EDLUAR (zolpidem) eszopiclone HETLIOZ (tasimelteon) <sup>CL*</sup> INTERMEZZO (zolpidem) LUNESTA (eszopiclone) ROZEREM (ramelteon) SILENOR (doxepin) SOMNOTE (chloral hydrate) SONATA (zaleplon) zaleplon zolpidem ER 6.25, 12.5 mg ZOLPIMIST (zolpidem)	Strengths of zolpidem that are non-preferred (6.25 and 12.5 mg) must be created by combining or splitting the preferred doses (5 and 10 mg) of zolpidem, if appropriate.  For treatment naïve female patients, zolpidem and zolpidem ER maximum dosages will be limited to 5 mg and 6.25 mg respectively per day.  *Full PA criteria may be found on the PA Criteria page by clicking the hyperlink.



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS			
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
SKELETAL MUSCLE RELAXANTS	P		
CLASS PA CRITERIA: See below for individual s	sub-class criteria.		
	ACUTE MUSCULOSKELETAL RELAXANT	AGENTS	
chlorzoxazone cyclobenzaprine IR 5, 10 mg methocarbamol	AMRIX (cyclobenzaprine) carisoprodol* carisoprodol/ASA* carisoprodol/ASA/codeine* cyclobenzaprine ER cyclobenzaprine IR 7.5 mg FEXMID (cyclobenzaprine) FLEXERIL (cyclobenzaprine) LORZONE (chlorzoxazone) metaxalone orphenadrine orphenadrine ER PARAFON FORTE (chlorzoxazone) ROBAXIN (methocarbamol) SKELAXIN (metaxalone) SOMA (carisoprodol)	Non-preferred agents require thirty (30) day trials of each preferred agent before they will be approved, unless one (1) of the exceptions on the PA form is present, with the exception of carisoprodol.  *Carisoprodol requires thirty (30) day trials of each of the preferred acute musculoskeletal relaxants and Skelaxin before it will be approved.	
	USCULOSKELETAL RELAXANT AGENTS USED F		
baclofen tizanidine tablets	DANTRIUM (dantrolene) dantrolene tizanidine capsules ZANAFLEX (tizanidine)	Non-preferred agents require thirty (30) day trials of each preferred agent before they will be approved, unless one (1) of the exceptions on the PA form is present.	
STEROIDS, TOPICAL			
CLASS PA CRITERIA: Non-preferred agents require five (5) day trials of one (1) form of EACH preferred unique active ingredient in the corresponding potency group before they will be approved, unless one (1) of the exceptions on the PA form is present.			
VERY HIGH & HIGH POTENCY			
betamethasone dipropionate cream betamethasone valerate cream betamethasone valerate lotion betamethasone valerate oint clobetasol propionate cream/gel/ointment/solution clobetasol emollient CLODAN SHAMPOO (clobetasol propionate) fluocinonide gel triamcinolone acetonide cream, ointment	amcinonide APEXICON (diflorasone diacetate) APEXICON E (diflorasone diacetate) betamethasone dipropionate gel, lotion, ointment clobetasol lotion, shampoo clobetasol propionate foam CLOBEX (clobetasol propionate) CLODAN KIT (clobetasol propionate) CORMAX (clobetasol propionate) desoximetasone cream/gel/ointment		



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
triamcinolone acetonide lotion	diflorasone diacetate DIPROLENE (betamethasone dipropionate/propylene glycol) DIPROLENE AF (betamethasone dipropionate/propylene glycol) DIPROSONE (betamethasone dipropionate) fluocinonide cream fluocinonide ointment fluocinonide solution fluocinonide/emollient halcinonide HALAC (halobetasol propionate) halobetasol propionate HALOG (halcinonide) HALONATE (halobetasol propionate) KENALOG (triamcinolone acetonide) LIDEX (fluocinonide) LIDEX-E (fluocinonide) OLUX (clobetasol propionate/emollient) PSORCON (diflorasone diacetate) SERNIVO SPRAY (betamethasone dipropionate) TEMOVATE (clobetasol propionate) TEMOVATE-E (clobetasol propionate) TEMOVATE-E (clobetasol propionate) TOPICORT CREAM, GEL, OINTMENT (desoximetasone) TOPICORT SPRAY (desoximetasone) ULTRAVATE (halobetasol propionate / lactic acid) VANOS (fluocinonide)	
MEDIUM POTENCY		
fluticasone propionate cream, ointment mometasone furoate triamcinolone acetonide 0.025% and 0.1% cream	ARISTOCORT (triamcinolone) BETA-VAL (betamethasone valerate) betamethasone valerate foam CLODERM (clocortolone pivalate) clocortolone cream CORDRAN/CORDRAN SP (flurandrenolide) CUTIVATE (fluticasone propionate) DERMATOP (prednicarbate) ELOCON (mometasone furoate)	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
	fluocinolone acetonide cream, ointment, solution fluticasone propionate lotion hydrocortisone butyrate cream hydrocortisone butyrate ointment, solution hydrocortisone valerate LOCOID (hydrocortisone butyrate) LOCOID LIPOCREAM (hydrocortisone butyrate/emollient) LUXIQ (betamethasone valerate) MOMEXIN (mometasone) PANDEL (hydrocortisone probutate) prednicarbate TOPICORT LP (desoximetasone) TRIDERM (triamcinolone acetonide) WESTCORT (hydrocortisone valerate)	
	LOW POTENCY	
hydrocortisone acetate (Rx, OTC) hydrocortisone cream (Rx, OTC) hydrocortisone lotion OTC hydrocortisone solution OTC hydrocortisone-aloe cream OTC hydrocortisone-aloe ointment OTC	ACLOVATE (alclometasone dipropionate) alclometasone dipropionate AQUA GLYCOLIC HC (hydrocortisone) CAPEX (fluocinolone acetonide) DERMA-SMOOTHE FS (fluocinolone acetonide) DESONATE (desonide) desonide cream, ointment desonide lotion DESOWEN (desonide) fluocinolone oil hydrocortisone/mineral oil/petrolatum hydrocortisone acetate/urea hydrocortisone lotion hydrocortisone/aloe gel LOKARA (desonide) PEDIADERM HC (hydrocortisone) PEDIADERM TA (hydrocortisone) SCALPICIN OTC (hydrocortisone) SYNALAR (fluocinolone) TEXACORT (hydrocortisone) TRIDESILON CREAM (desonide) VERDESO (desonide)	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS			
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
STIMULANTS AND RELATED AGENTS  CLASS PA CRITERIA: A PA is required for adults eighteen (18) years of age or older.  PLEASE NOTE: Requests for IR + ER combination therapy must be for the same active ingredient in the same salt form, if available.			
exceptions on the PA form is present. NOTE: Non-	preferred agents will NOT be "grandfathered" for adult they will be required to switch to a preferred agent.	and with a similar duration of effect, unless one (1) of the ults. Children under the age of 18 may continue their current	
	AMPHETAMINES		
ADZENYS XR ODT (amphetamine) amphetamine salt combination IR dextroamphetamine ER dextroamphetamine IR PROCENTRA solution (dextroamphetamine) VYVANSE CHEWABLE (lisdexamfetamine) VYVANSE CAPSULE (lisdexamfetamine)	ADDERALL (amphetamine salt combination) ADDERALL XR* (amphetamine salt combination) amphetamine salt combination ER DESOXYN (methamphetamine) DEXEDRINE ER (dextroamphetamine) DEXEDRINE IR (dextroamphetamine) dextroamphetamine solution DYANAVEL XR SUSP (amphetamine) EVEKEO (amphetamine) methamphetamine	In addition to the Class Criteria: Thirty (30) day trials of at least three (3) antidepressants are required before amphetamines will be authorized for depression.  *Adderall XR is preferred over its generic equivalents.  **Mydayis requires a 30-day trial of at least one long-acting preferred agent in this subclass and a trial of Adderall XR.	
	MYDAYIS (dextroamphetamine/amphetamine salt)** ZENZEDI (dextroamphetamine)  NON-AMPHETAMINE		
APTENSIO XR (methylphenidate)	clonidine ER*	*Kapvay/clonidine ER will be authorized only after fourteen (14)	
armodafinil <sup>CL</sup> atomoxetine clonidine IR  COTEMPLA XR ODT (methylphenidate) DAYTRANA (methylphenidate) dexmethylphenidate IR FOCALIN XR (dexmethylphenidate) guanfacine ER guanfacine IR METADATE CD (methylphenidate) discontinued by labeler METHYLIN SOLUTION (methylphenidate) methylphenidate IR modafinil <sup>CL</sup> QUILLICHEW ER (methylphenidate)	CONCERTA (methylphenidate) dexmethylphenidate XR FOCALIN IR (dexmethylphenidate) INTUNIV (guanfacine extended-release) KAPVAY (clonidine extended-release)* methylphenidate CD methylphenidate chewable tablets, solution methylphenidate ER methylphenidate ER (generic CONCERTA) methylphenidate LA NUVIGIL (armodafinil) PROVIGIL (modafinil) QUILLIVANT XR (methylphenidate) RITALIN (methylphenidate) RITALIN LA (methylphenidate) STRATTERA (atomoxetine)**	day trials of at least one (1) preferred product from both the amphetamine and non-amphetamine class. These trials must include a fourteen (14) day trial of clonidine IR unless one (1) of the exceptions on the PA form is present.  NOTE: In cases of a diagnosis of Tourette's syndrome, tics, autism or disorders included in the autism spectrum, Kapvay will only require a fourteen (14) day trial of clonidine IR for approval.  **Strattera is limited to a maximum of 100 mg per day.	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS				
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA		
TETRACYCLINES				
<b>CLASS PA CRITERIA:</b> Non-preferred agents re PA form is present.	CLASS PA CRITERIA: Non-preferred agents require ten (10) day trials of each preferred agent before they will be approved, unless one (1) of the exceptions on the PA form is present.			
doxycycline hyclate capsules doxycycline hyclate 50, 100 mg tablets doxycycline monohydrate 50, 100 mg capsules minocycline capsules tetracycline	ADOXA (doxycycline monohydrate) demeclocycline* DORYX (doxycycline hyclate) doxycycline hyclate 75, 150 mg tablets doxycycline hyclate tablet DR doxycycline monohydrate 40, 75, 150 mg capsule doxycycline monohydrate tablet doxycycline monohydrate suspension DYNACIN (minocycline) MINOCIN (minocycline) minocycline ER capsules minocycline tablets MONODOX (doxycycline monohydrate) MORGIDOX KIT (doxycycline) ORACEA (doxycycline monohydrate) SOLODYN (minocycline) VIBRAMYCIN CAPSULES, SUSPENSION, SYRUP (doxycycline)	*Demeclocycline will be authorized for conditions caused by susceptible strains of organisms designated in the product information supplied by the manufacturer. A C&S report must accompany this request.  Demeclocycline will also be authorized for SIADH.		
ULCERATIVE COLITIS AGENTSAP				
CLASS PA CRITERIA: Non-preferred agents require thirty (30) day trials of each preferred dosage form or chemical entity before the corresponding non-preferred agent of that dosage form or chemical entity will be approved, unless one (1) of the exceptions on the PA form is present.				
ORAL				
APRISO (mesalamine) balsalazide	ASACOL HD (mesalamine) AZULFIDINE (sulfasalazine)			

	ORAL	
APRISO (mesalamine) balsalazide sulfasalazine	ASACOL HD (mesalamine) AZULFIDINE (sulfasalazine) COLAZAL (balsalazide) DELZICOL (mesalamine) DIPENTUM (olsalazine) GIAZO (balsalazide) LIALDA (mesalamine) mesalamine PENTASA (mesalamine) 250 mg PENTASA (mesalamine) 500 mg UCERIS (budesonide)	



EFFECTIVE 07/01/2018 Version 2018.3e

THERAPEUTIC DRUG CLASS			
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA	
	RECTAL		
CANASA (mesalamine) mesalamine	DELZICOL DR (mesalamine) mesalamine kit ROWASA (mesalamine) SF ROWASA (mesalamine) UCERIS (budesonide)		
VASODILATORS, CORONARY			
CLASS PA CRITERIA: Non-preferred agents require thirty (30) day trials of each preferred dosage form before they will be approved, unless one (1) of the exceptions on the PA form is present.			
SUBLINGUAL NITROGLYCERIN			
nitroglycerin spray (generic NITROLINGUAL) nitroglycerin sublingual NITROSTAT SUBLINGUAL (nitroglycerin)	GONITRO SPRAY POWDER (nitroglycerin) nitroglycerin spray (generic NITROMIST) NITROLINGUAL SPRAY (nitroglycerin) NITROMIST (nitroglycerin)		